

# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/GB05/000779

International filing date: 03 March 2005 (03.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: GB  
Number: 0405055.5  
Filing date: 05 March 2004 (05.03.2004)

Date of receipt at the International Bureau: 21 April 2005 (21.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland  
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



INVESTOR IN PEOPLE

The Patent Office  
Concept House  
Cardiff Road  
Newport  
South Wales  
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.


Signed

Dated 21 March 2005



## Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)



The Patent Office

Cardiff Road  
Newport  
South Wales  
NP9 1RH

1. Your reference

05 MAR 2004

2. Patent application number

(The Patent Office will fill in this part)

0405055.5

3. Full name, address and postcode of the or of each applicant (*underline all surnames*)

Eisai London Research Laboratories Limited  
Bernard Katz Building  
University College London  
Gower Street  
London WC1E 6BT

Patents ADP number (*if you know it*)

If the applicant is a corporate body, give the country/state of its incorporation

6347256001

4. Title of the invention

JNK Inhibitors

5. Name of your agent (*if you have one*)

"Address for service" in the United Kingdom to which all correspondence should be sent (*including the postcode*)

Kilburn & Strode  
20 Red Lion Street  
London  
WC1R 4PJ

Patents ADP number (*if you know it*)

125001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (*if you know it*) the or each application number

Country

Priority application number  
(*if you know it*)

Date of filing  
(*day / month / year*)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing  
(*day / month / year*)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (*Answer 'Yes' if:*

- a) any applicant named in part 3 is not an inventor, or
  - b) there is an inventor who is not named as an applicant, or
  - c) any named applicant is a corporate body.
- See note (d))

YES



## Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description

Claim(s)

Abstract

Drawing(s)

54 53  
32 33  
52

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (*Patents Form 7/77*)

Request for preliminary examination and search (*Patents Form 9/77*)

Request for substantive examination (*Patents Form 10/77*)

Any other documents  
(*please specify*)

11. I/We request the grant of a patent on the basis of this application.

Signature

Date 5 March 2004

Kubun & Stode

12. Name and daytime telephone number of person to contact in the United Kingdom

Kristina Cornish  
Tel: 020 7539 4200

### Warning

*After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.*

### Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.*
- Write your answers in capital letters using black ink or you may type them.*
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.*
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.*
- Once you have filled in the form you must remember to sign and date it.*
- For details of the fee and ways to pay please contact the Patent Office.*

## JNK Inhibitors

The present invention relates to novel compounds, their use in the inhibition of c-Jun N-terminal kinases, their use in medicine and particularly in the prevention and/or treatment of neurodegenerative disorders related to apoptosis and/or inflammation. The invention also provides processes for manufacture of said compounds, compositions containing them and processes for manufacturing such compositions.

c-Jun N-terminal kinases (hereinafter referred to as "JNKs") are members of the mitogen-activated protein kinase (MAPK) family. JNKs are involved in response to various stimuli, including proinflammatory cytokines and environmental stress. JNKs, and JNK3 in particular, play an important role during apoptotic death of cells and therefore have been implicated in various disorders including stroke, traumatic brain injury and other neurodegenerative diseases such as Parkinson disease, Alzheimer disease and others. Since JNK activity is a physiological regulator of AP-1 transcriptional activity, JNK inhibitors are expected to reduce inflammatory response.

Apoptosis is a form of cell death in which the cell actively participates in its own destruction in a process involving a characteristic series of biochemical and morphological changes, which are regulated by specific cell death genes. The apoptotic cell death is a process that has been observed in the developing mammalian nervous system. In mice, the inactivation by homologous recombination of genes that encode proteins that promote apoptosis, such as the caspase-3 or the Bax protein, prevents developmental neuronal cell death. The destruction of genes that encode cell death suppressors such as Bcl-x, leads to enhanced neuronal cell death. There is increasing evidence that apoptosis plays an important role in the pathology of acute and chronic neurodegenerative

diseases. For example, in transgenic mice overexpressing the anti-apoptotic Bcl-2 protein in the nervous system there is a decrease in infarct volume following cerebral ischemia. Similarly, injection of the caspase inhibitor BAF reduces neuronal cell death following hypoxia/ischaemia in neonatal rats.

5 Another example is spinal muscular atrophy (a motor neuron disease) where loss of function mutations in the SMN gene is associated with the disease. Recent data has shown that the wild type SMN protein binds to Bcl-2 and co-operates with it to inhibit apoptosis. These results suggest that inhibitors of neuronal apoptosis could be beneficial in the treatment of human  
10 neurodegenerative diseases. There is increasing evidence that neuronal apoptosis is an important pathological feature of stroke, traumatic brain injury and other neurodegenerative diseases. Therefore, pharmacotherapy using inhibitors of neuronal apoptosis may provide a therapeutic benefit in neurodegenerative conditions.

15

A number of groups have studied the mechanisms of neuronal cell death using in vitro cell culture systems and the results suggest that in some systems the transcription factor c-Jun is activated by the removal of survival signals and promotes cell death.

20

Antibodies specific for c-Jun protected NGF-deprived rat sympathetic neurones from apoptosis. Analogous neuroprotection due to expression of a c-Jun dominant negative mutant has been demonstrated, whereas overexpression of wild type c-Jun protein was sufficient to induce apoptosis in the presence of  
25 NGF. Estus and co-workers recently showed that an increase in c-Jun RNA levels occurs in cortical neurones undergoing apoptosis after treatment with  $\beta$ -amyloid peptide. It has also been shown that c-Jun is required for apoptosis in cerebellar granule neurones deprived of survival signals.

c-Jun is activated by JNKs, which phosphorylate its transcriptional activation domain. In humans there are three JNK genes: JNK1, JNK2 and JNK3. The RNAs encoding JNK1 and JNK2 are expressed in many tissues, including the brain, but JNK3 is restricted to the nervous system and to a smaller extent the  
5 heart and testes.

JNKs are strongly activated in cellular responses to various stresses such as UV radiation, heat shock, osmotic shock, DNA-damaging agents, and proinflammatory cytokines such as  $\text{TNF}\alpha$ ,  $\text{IL-1}\beta$  and others. Upstream  
10 regulators of the JNK pathway include kinases such as SEK1, MKK7 and MEKK1. There is evidence that Jun kinase activity is required for neuronal apoptosis in vitro. Overexpression of MEKK1 in sympathetic neurones increased c-Jun protein levels and phosphorylation and induced apoptosis in the presence of NGF indicating that activation of the Jun kinase pathway can  
15 trigger neuronal cell death. The Jun kinase pathway has been shown to be necessary for the death of differentiated PC12 cells deprived of NGF. Furthermore, compound CEP-1347, which inhibits the c-Jun pathway (upstream of Jun kinase), protects motor neurones against cell death induced by survival factor withdrawal.

20

In JNK3 homozygous (-/-) knockout mice, epileptic seizures and death of hippocampal CA3 neurones induced by injection of kainic acid is blocked. This indicates that JNK3 is involved in certain forms of neuronal cell death in vivo. It is also a critical component of GluR6-mediated excitotoxicity. Furthermore,  
25 JNK3 (-/-) mice appear to develop normally and are viable suggesting that JNK3 is not essential for development or viability.


Strong nuclear JNK3 immunoreactivity in the brain CA1 neurones of patients with acute hypoxia suggests that JNK3 is involved in hypoxia-related

neurodegeneration. Transient hypoxia may also trigger apoptosis through JNK signaling pathway in developing brain neurones.

Furthermore, JNK3 immunoreactivity is colocalized with Alzheimer disease-  
5 affected neurones. Moreover JNK3 is related to neurofibrillary pathology of Alzheimer disease. In particular, JNK3 induces robust phosphorylation of amyloid precursor protein (APP) thus affecting its metabolism in disease state.


The present inventors have provided compounds, which are inhibitors of c-Jun  
10 N-terminal kinases.

The first aspect of the invention therefore relates to a compound of formula (I) as illustrated below:



(I)

15 wherein R¹ is an optionally substituted C<sub>3-12</sub> carbocyclyl or C<sub>3-12</sub> heterocyclyl group or a group of formula (II)



(II)

wherein X is NR<sup>3</sup>, O, S or (CR<sup>22</sup>R<sup>22</sup>)<sub>n</sub>, Y is absent or is NR<sup>23</sup>, O, or (CR<sup>23</sup>R<sup>23</sup>)<sub>n</sub>,  
20 R² is optionally substituted C<sub>1-12</sub> alkyl, C<sub>2-12</sub> alkenyl, C<sub>2-12</sub> alkynyl, C<sub>3-12</sub> carbocyclyl or C<sub>3-12</sub> heterocyclyl, and R⁴ is an optionally substituted five or six membered heterocyclyl group or an optionally substituted six membered carbocyclyl group;



wherein the optionally substituted carbocyclyl or heterocyclyl group of  $R^1$  is optionally fused to a partially saturated, unsaturated or fully saturated five to seven membered ring containing zero to three heteroatoms, and each substitutable carbon atom in  $R^1$ , including the optional fused ring, is optionally and independently substituted by one or more of halogen,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl, halo $C_{1-12}$ alkyl,  $C_{3-12}$  carbocyclyl,  $C_{3-12}$  heterocyclyl,  $(CH_2)_nOR^5$ ,  $(CH_2)_nNR^5_2(CH_2)_nSR^5$ ,  $OR^5$ ,  $SR^5$ ,  $NO_2$ ,  $CN$ ,  $NR^5_2$ ,  $NR^5COR^5$ ,  $NR^5CONR^5_2$ ,  $NR^5COR^5$ ,  $NR^5CO_2R^5$ ,  $CO_2R^5$ ,  $COR^5$ ,  $CONR^5_2$ ,  $S(O)_2R^5$ ,  $SONR^5_2$ ,  $S(O)R^5$ ,  $SO_2NR^5_2$ , or  $NR^5S(O)_2R^5$  wherein the  $C_{1-12}$  alkyl group optionally contains one or more insertions selected from  $-O-$ ,  $-N(R^5)-$ ,  $-S-$ ,  $-S(O)-$  and  $-S(O_2)-$ ; and each saturated carbon in the optional fused ring is further optionally and independently substituted by  $=O$ ,  $=S$ ,  $NNR^6_2$ ,  $=N-OR^6$ ,  $=NNR^6COR^6$ ,  $=NNR^6CO_2R^6$ ,  $=NNSO_2R^6$ , or  $=NR^6$ ; and each substitutable nitrogen atom in  $R^1$  is optionally substituted by  $R^7$ ,  $COR^7$ ,  $SO_2R^7$  or  $CO_2R^7$ ;

wherein  $n$  is 1 to 6, preferably  $n$  is 1, 2 or 3;

wherein  $R^5$  is hydrogen,  $C_{1-12}$  alkyl,  $C_{3-12}$  carbocyclyl or  $C_{3-12}$  heterocyclyl, optionally substituted by one or more of  $C_{1-6}$  alkyl,  $C_{3-12}$  carbocyclyl,  $C_{3-12}$  heterocyclyl, halogen,  $C_{1-6}$  haloalkyl,  $OR^8$ ,  $SR^8$ ,  $NO_2$ ,  $CN$ ,  $NR^8R^8$ ,  $NR^8COR^8$ ,  $NR^8CONR^8R^8$ ,  $NR^8COR^8$ ,  $NR^8CO_2R^8$ ,  $CO_2R^8$ ,  $COR^8$ ,  $CONR^8_2$ ,  $S(O)_2R^8$ ,  $SONR^8_2$ ,  $S(O)R^8$ ,  $SO_2NR^8R^8$ ,  $NR^8S(O)_2R^8$ , wherein the  $C_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-O-$ ,  $-N(R^8)-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^8$  may be the same or different and is as defined below;

25

wherein two  $R^5$  in  $NR^5_2$  may optionally form a partially saturated, unsaturated or fully saturated three to seven membered ring containing one to three heteroatoms, optionally and independently substituted by one or more of  $C_{1-6}$  alkyl, halogen,  $C_{1-6}$  haloalkyl,  $OR^8$ ,  $SR^8$ ,  $NO_2$ ,  $CN$ ,  $NR^8R^8$ ,  $NR^8COR^8$ ,

$\text{NR}^8\text{CONR}^8\text{R}^8$ ,  $\text{NR}^8\text{COR}^8$ ,  $\text{NR}^8\text{CO}_2\text{R}^8$ ,  $\text{CO}_2\text{R}^8$ ,  $\text{COR}^8$ ,  $\text{CONR}^8_2$ ,  $\text{S(O)}_2\text{R}^8$ ,  
 $\text{SONR}^8_2$ ,  $\text{S(O)R}^8$ ,  $\text{SO}_2\text{NR}^8\text{R}^8$ ,  $\text{NR}^8\text{S(O)}_2\text{R}^8$ ,

wherein the  $\text{C}_{1-6}$  alkyl group optionally incorporates one or two insertions  
 selected from the group consisting of  $-\text{O}-$ ,  $-\text{N(R}^8)-$ ,  $-\text{S(O)}-$  and  $-\text{S(O)}_2-$ , wherein  
 5 each  $\text{R}^8$  may be the same or different and is as defined below;

wherein  $\text{R}^6$  is hydrogen,  $\text{C}_{1-12}$  alkyl,  $\text{C}_{3-12}$  carbocyclyl or  $\text{C}_{3-12}$  heterocyclyl,  
 optionally substituted by one or more of  $\text{C}_{1-6}$  alkyl, halogen,  $\text{C}_{1-6}$  haloalkyl,  
 $\text{OR}^8$ ,  $\text{SR}^8$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{NR}^8\text{R}^8$ ,  $\text{NR}^8\text{COR}^8$ ,  $\text{NR}^8\text{CONR}^8\text{R}^8$ ,  $\text{NR}^8\text{COR}^8$ ,  $\text{NR}^8\text{CO}_2\text{R}^8$ ,  
 10  $\text{CO}_2\text{R}^8$ ,  $\text{COR}^8$ ,  $\text{CONR}^8_2$ ,  $\text{S(O)}_2\text{R}^8$ ,  $\text{S(O)R}^8$ ,  $\text{SO}_2\text{NR}^8\text{R}^8$ ,  $\text{NR}^8\text{S(O)}_2\text{R}^8$ , wherein  
 the  $\text{C}_{1-12}$  alkyl group optionally incorporates one or two insertions selected from  
 the group consisting of  $-\text{O}-$ ,  $-\text{N(R}^8)-$ ,  $-\text{S(O)}-$  and  $-\text{S(O)}_2-$ , wherein each  $\text{R}^8$  may  
 be the same or different and is as defined below;

15 wherein  $\text{R}^7$  is hydrogen,  $\text{C}_{6-12}$  aryl,  $\text{C}_{1-6}$  alkyl or  $\text{C}_{1-6}$  haloalkyl;

wherein  $\text{R}^8$  is hydrogen,  $\text{C}_{1-6}$  alkyl, or  $\text{C}_{1-6}$  haloalkyl;

$\text{Y}$  is absent or is  $\text{NR}^{23}$ ,  $\text{O}$ , or  $(\text{CR}^{23}\text{R}^{23})_n$ , wherein each  $\text{R}^{23}$  may be the same or  
 20 different and is  $\text{H}$ ,  $\text{C}_{1-4}$  alkyl,  $\text{C}_{1-4}$  alkoxy or  $\text{C}_{1-4}$  haloalkyl;  
 and  $n$  is 1 to 6, preferably  $n$  is 1, 2, 3 or 4.

$\text{R}^2$  is  $\text{C}_{1-12}$  alkyl,  $\text{C}_{2-12}$  alkenyl,  $\text{C}_{2-12}$  alkynyl,  $\text{C}_{3-12}$  carbocyclyl or  $\text{C}_{3-12}$   
 heterocyclyl, each of which is optionally substituted, wherein;

25 the optionally substituted carbocyclyl or heterocyclyl group is optionally fused  
 to one to three unsaturated, partially unsaturated or fully saturated five to seven  
 membered rings containing zero to three heteroatoms;

each substitutable carbon atom in  $\text{R}^2$ , including the optional fused ring, is  
 optionally and independently substituted by one or more of  $\text{C}_{1-12}$  alkyl,  $\text{C}_{3-12}$



- cycloalkyl, C<sub>3-12</sub> heterocycloalkyl, C<sub>3-12</sub> aryl, C<sub>3-12</sub> heteroaryl halogen, C<sub>1-12</sub> haloalkyl, OR<sup>9</sup>, SR<sup>9</sup>, NO<sub>2</sub>, CN, NR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>COR<sup>9</sup>, NR<sup>9</sup>CONR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>COR<sup>9</sup>, NR<sup>9</sup>CO<sub>2</sub>R<sup>9</sup>, CO<sub>2</sub>R<sup>9</sup>, COR<sup>9</sup>, CONR<sup>9</sup>R<sup>9</sup>, S(O)<sub>2</sub>R<sup>9</sup>, SONH<sub>2</sub>, S(O)R<sup>9</sup>, SO<sub>2</sub>NR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>S(O)<sub>2</sub>R<sup>9</sup>, wherein each R<sup>9</sup> may be the same or different and is as defined
- 5 below and wherein:
- the C<sub>1-12</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -C(O)-, -N(R<sup>9</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>9</sup> may be the same or different and is as defined above;
- the C<sub>1-12</sub> alkyl, C<sub>3-12</sub> cycloalkyl, C<sub>3-12</sub> heterocycloalkyl, C<sub>3-12</sub> aryl, or C<sub>3-12</sub> heteroaryl groups are optionally substituted by one or more of halogen, C<sub>1-12</sub> haloalkyl, OR<sup>9</sup>, SR<sup>9</sup>, NO<sub>2</sub>, CN, NR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>COR<sup>9</sup>, NR<sup>9</sup>CONR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>COR<sup>9</sup>, NR<sup>9</sup>CO<sub>2</sub>R<sup>9</sup>, CO<sub>2</sub>R<sup>9</sup>, COR<sup>9</sup>, CONR<sup>9</sup>R<sup>9</sup>, S(O)<sub>2</sub>R<sup>9</sup>, SONH<sub>2</sub>, S(O)R<sup>9</sup>, SO<sub>2</sub>NR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>S(O)<sub>2</sub>R<sup>9</sup>, wherein each R<sup>9</sup> may be the same or different and is as defined below; and
- 15 the C<sub>3-12</sub> cycloalkyl, C<sub>3-12</sub> heterocycloalkyl, C<sub>3-12</sub> aryl, or C<sub>3-12</sub> heteroaryl groups are optionally substituted by one or more C<sub>1-12</sub> alkyl groups;
- each saturated carbon in R<sup>2</sup>, including the optional fused ring, is further optionally and independently substituted by =O, =S, NNR<sup>9</sup>R<sup>9</sup>, =N-OR<sup>9</sup>, =NNHCOR<sup>9</sup>, =NNHCO<sub>2</sub>R<sup>9</sup>, =NNSO<sub>2</sub>R<sup>9</sup>, or =NR<sup>9</sup>, wherein each R<sup>9</sup> may be the
- 20 same or different and is as defined below; and
- each substitutable nitrogen atom in R<sup>2</sup> is optionally substituted by R<sup>10</sup>, COR<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup> or CO<sub>2</sub>R<sup>9</sup> wherein each R<sup>9</sup> and R<sup>10</sup> may be the same or different and is as defined below;
- wherein two R<sup>9</sup> in NR<sup>9</sup><sub>2</sub> may optionally form a partially saturated, unsaturated
- 25 or fully saturated three to seven membered ring containing one to three heteroatoms, optionally and independently substituted by one or more of C<sub>1-6</sub> alkyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>11</sup>, SR<sup>11</sup>, NO<sub>2</sub>, CN, NR<sup>11</sup>R<sup>11</sup>, NR<sup>11</sup>COR<sup>11</sup>, NR<sup>11</sup>CONR<sup>11</sup>R<sup>11</sup>, NR<sup>11</sup>COR<sup>11</sup>, NR<sup>11</sup>CO<sub>2</sub>R<sup>11</sup>, CO<sub>2</sub>R<sup>11</sup>, COR<sup>11</sup>, CONR<sup>11</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>11</sup>, SONR<sup>11</sup><sub>2</sub>, S(O)R<sup>11</sup>, SO<sub>2</sub>NR<sup>11</sup>R<sup>11</sup>, NR<sup>11</sup>S(O)<sub>2</sub>R<sup>11</sup>,

wherein the C<sub>1-6</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -N(R<sup>11</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>11</sup> may be the same or different and is as defined below; wherein R<sup>11</sup> is hydrogen, C<sub>1-6</sub> alkyl, or C<sub>1-6</sub> haloalkyl;

5

wherein R<sup>9</sup> is hydrogen, C<sub>1-12</sub> alkyl or C<sub>3-12</sub> aryl, optionally substituted by one or more of C<sub>1-4</sub> alkyl, halogen, C<sub>1-4</sub> haloalkyl, OR<sup>12</sup>, SR<sup>12</sup>, NO<sub>2</sub>, CN, NR<sup>12</sup>R<sup>12</sup>, NR<sup>12</sup>COR<sup>12</sup>, NR<sup>12</sup>CONR<sup>12</sup>R<sup>12</sup>, NR<sup>12</sup>COR<sup>12</sup>, NR<sup>12</sup>CO<sub>2</sub>R<sup>12</sup>, CO<sub>2</sub>R<sup>12</sup>, COR<sup>12</sup>, CONR<sup>12</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>12</sup>, SONH<sub>2</sub>, S(O)R<sup>12</sup>, SO<sub>2</sub>NR<sup>12</sup>R<sup>12</sup>, NR<sup>12</sup>S(O)<sub>2</sub>R<sup>12</sup>, wherein  
10 the C<sub>1-12</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -N(R<sup>12</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>12</sup> may be the same or different and is as defined below;

wherein R<sup>10</sup> is C<sub>1-12</sub> alkyl or C<sub>3-12</sub> aryl, optionally substituted by one or more of  
15 C<sub>1-4</sub> alkyl, halogen, C<sub>1-4</sub> haloalkyl, OR<sup>12</sup>, SR<sup>12</sup>, NO<sub>2</sub>, CN, NR<sup>12</sup>R<sup>12</sup>, NR<sup>12</sup>COR<sup>12</sup>, NR<sup>12</sup>CONR<sup>12</sup>R<sup>12</sup>, NR<sup>12</sup>COR<sup>12</sup>, NR<sup>12</sup>CO<sub>2</sub>R<sup>12</sup>, CO<sub>2</sub>R<sup>12</sup>, COR<sup>12</sup>, CONR<sup>12</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>12</sup>, SONH<sub>2</sub>, S(O)R<sup>12</sup>, SO<sub>2</sub>NR<sup>12</sup>R<sup>12</sup>, NR<sup>12</sup>S(O)<sub>2</sub>R<sup>12</sup>, wherein the C<sub>1-12</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -N(R<sup>12</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>12</sup> may be the  
20 same or different and is as defined below;

wherein R<sup>12</sup> is hydrogen, C<sub>1-4</sub> alkyl, or C<sub>1-4</sub> haloalkyl;

X is NR<sup>3</sup>; O, S or (CR<sup>22</sup>R<sup>22</sup>)<sub>n</sub> wherein R<sup>22</sup> is independently one or more of  
25 halogen, C<sub>1-12</sub> alkyl, C<sub>2-12</sub> alkenyl, C<sub>2-12</sub> alkynyl, C<sub>1-12</sub> haloalkyl, C<sub>6-12</sub> carbocyclyl, C<sub>5-12</sub> heterocyclyl, (CH<sub>2</sub>)<sub>n</sub>OR<sup>5</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>5</sup><sub>2</sub>, OR<sup>5</sup>, SR<sup>5</sup>, NO<sub>2</sub>, CN, NR<sup>5</sup><sub>2</sub>, NR<sup>5</sup>COR<sup>5</sup>, NR<sup>5</sup>CONR<sup>5</sup><sub>2</sub>, NR<sup>5</sup>COR<sup>5</sup>, NR<sup>5</sup>CO<sub>2</sub>R<sup>5</sup>, CO<sub>2</sub>R<sup>5</sup>, COR<sup>5</sup>, CONR<sup>5</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>5</sup>, SONR<sup>5</sup><sub>2</sub>, S(O)R<sup>5</sup>, SO<sub>2</sub>NR<sup>5</sup><sub>2</sub>, or NR<sup>5</sup>S(O)<sub>2</sub>R<sup>5</sup> wherein each R<sup>5</sup> may be the same or different and is as defined above; and

wherein  $n$  is 1 to 6, preferably  $n$  is 1, 2, 3 or 4;

wherein  $R^3$  is a lone electron pair, hydrogen,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{3-12}$  carbocyclyl or  $C_{3-12}$  heterocyclyl, each of which is optionally  
5 substituted, wherein:

the optionally substituted carbocyclyl or heterocyclyl group is optionally fused to one to three unsaturated, partially unsaturated or fully saturated five to seven membered rings containing zero to three heteroatoms,

each substitutable carbon atom in  $R^3$ , including the optional fused ring, is  
10 optionally and independently substituted by one or more of  $C_{1-12}$  alkyl,  $C_{3-12}$  cycloalkyl,  $C_{3-12}$  heterocycloalkyl,  $C_{3-12}$  aryl,  $C_{3-12}$  heteroaryl halogen,  $C_{1-12}$  haloalkyl,  $OR^{13}$ ,  $SR^{13}$ ,  $NO_2$ ,  $CN$ ,  $NR^{13}R^{13}$ ,  $NR^{13}COR^{13}$ ,  $NR^{13}CONR^{13}R^{13}$ ,  $NR^{13}COR^{13}$ ,  $NR^{13}CO_2R^{13}$ ,  $CO_2R^{13}$ ,  $COR^{13}$ ,  $CONR^{13}R^{13}$ ,  $S(O)_2R^{13}$ ,  $SONH_2$ ,  $S(O)R^{13}$ ,  $SO_2NR^{13}R^{13}$ ,  $NR^{13}S(O)_2R^{13}$ , wherein each  $R^{13}$  may be the same or  
15 different and is as defined above and wherein:

the  $C_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-O-$ ,  $-C(O)-$ ,  $-N(R^{13})-$ ,  $-S(O)-$  and  $-S(O)_2-$ , wherein each  $R^{13}$  may be the same or different and is as defined above;

the  $C_{1-12}$  alkyl,  $C_{3-12}$  cycloalkyl,  $C_{3-12}$  heterocycloalkyl,  $C_{3-12}$  aryl, or  $C_{3-12}$   
20 heteroaryl groups are optionally substituted by one or more of halogen,  $C_{1-12}$  haloalkyl,  $OR^{13}$ ,  $SR^{13}$ ,  $NO_2$ ,  $CN$ ,  $NR^{13}R^{13}$ ,  $NR^{13}COR^{13}$ ,  $NR^{13}CONR^{13}R^{13}$ ,  $NR^{13}COR^{13}$ ,  $NR^{13}CO_2R^{13}$ ,  $CO_2R^{13}$ ,  $COR^{13}$ ,  $CONR^{13}R^{13}$ ,  $S(O)_2R^{13}$ ,  $SONH_2$ ,  $S(O)R^{13}$ ,  $SO_2NR^{13}R^{13}$ ,  $NR^{13}S(O)_2R^{13}$ , wherein each  $R^{13}$  may be the same or different and is as defined below; and

25 the  $C_{3-12}$  cycloalkyl,  $C_{3-12}$  heterocycloalkyl,  $C_{3-12}$  aryl, or  $C_{3-12}$  heteroaryl groups are optionally substituted by one or more  $C_{1-12}$  alkyl groups;

each saturated carbon in  $R^2$ , including the optional fused ring, is further optionally and independently substituted by  $=O$ ,  $=S$ ,  $NNR^{13}R^{13}$ ,  $=N-OR^{13}$ ,

$=\text{NNHCOR}^{13}$ ,  $=\text{NNHCO}_2\text{R}^{13}$ ,  $=\text{NNSO}_2\text{R}^{13}$ , or  $=\text{NR}^{13}$ , wherein each  $\text{R}^{13}$  may be the same or different and is as defined below; and

each substitutable nitrogen atom in  $\text{R}^3$  is optionally substituted by  $\text{R}^{14}$ ,  $\text{COR}^{13}$ ,  $\text{SO}_2\text{R}^{13}$  or  $\text{CO}_2\text{R}^{13}$  wherein each  $\text{R}^{13}$  and  $\text{R}^{14}$  may be the same or different and is as defined below;

5 wherein two  $\text{R}^{13}$  in  $\text{NR}^{13}_2$  may optionally form a partially saturated, unsaturated or fully saturated three to seven membered ring containing one to three heteroatoms, optionally and independently substituted by one or more of  $\text{C}_{1-6}$  alkyl, halogen,  $\text{C}_{1-6}$  haloalkyl,  $\text{OR}^{15}$ ,  $\text{SR}^{15}$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{NR}^{15}\text{R}^{15}$ ,  $\text{NR}^{15}\text{COR}^{15}$ ,  
 10  $\text{NR}^{15}\text{CONR}^{15}\text{R}^{15}$ ,  $\text{NR}^{15}\text{COR}^{15}$ ,  $\text{NR}^{15}\text{CO}_2\text{R}^{15}$ ,  $\text{CO}_2\text{R}^{15}$ ,  $\text{COR}^{15}$ ,  $\text{CONR}^{15}_2$ ,  $\text{S(O)}_2\text{R}^{15}$ ,  $\text{SONR}^{15}_2$ ,  $\text{S(O)R}^{15}$ ,  $\text{SO}_2\text{NR}^{15}\text{R}^{15}$ ,  $\text{NR}^{15}\text{S(O)}_2\text{R}^{15}$ ,

wherein the  $\text{C}_{1-6}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-\text{O}-$ ,  $-\text{N}(\text{R}^{15})-$ ,  $-\text{S(O)}-$  and  $-\text{S(O)}_2-$ , wherein each  $\text{R}^{15}$  may be the same or different and is as defined below;

15 wherein  $\text{R}^{15}$  is hydrogen,  $\text{C}_{1-6}$  alkyl, or  $\text{C}_{1-6}$  haloalkyl;

wherein  $\text{R}^{13}$  is hydrogen,  $\text{C}_{1-12}$  alkyl or  $\text{C}_{3-12}$  aryl, optionally substituted by one or more of  $\text{C}_{1-4}$  alkyl, halogen,  $\text{C}_{1-4}$  haloalkyl,  $\text{OR}^{16}$ ,  $\text{SR}^{16}$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{NR}^{16}\text{R}^{16}$ ,  $\text{NR}^{16}\text{COR}^{16}$ ,  $\text{NR}^{16}\text{CONR}^{16}\text{R}^{16}$ ,  $\text{NR}^{16}\text{COR}^{16}$ ,  $\text{NR}^{16}\text{CO}_2\text{R}^{16}$ ,  $\text{CO}_2\text{R}^{16}$ ,  $\text{COR}^{16}$ ,  
 20  $\text{CONR}^{16}_2$ ,  $\text{S(O)}_2\text{R}^{16}$ ,  $\text{SONH}_2$ ,  $\text{S(O)R}^{16}$ ,  $\text{SO}_2\text{NR}^{16}\text{R}^{16}$ ,  $\text{NR}^{16}\text{S(O)}_2\text{R}^{16}$ , wherein the  $\text{C}_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-\text{O}-$ ,  $-\text{N}(\text{R}^{16})-$ ,  $-\text{S(O)}-$  and  $-\text{S(O)}_2-$ , wherein each  $\text{R}^{16}$  may be the same or different and is as defined below;

25 wherein  $\text{R}^{14}$  is  $\text{C}_{1-12}$  alkyl or  $\text{C}_{3-12}$  aryl, optionally substituted by one or more of  $\text{C}_{1-4}$  alkyl, halogen,  $\text{C}_{1-4}$  haloalkyl,  $\text{OR}^{16}$ ,  $\text{SR}^{16}$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{NR}^{16}\text{R}^{16}$ ,  $\text{NR}^{16}\text{COR}^{16}$ ,  $\text{NR}^{16}\text{CONR}^{16}\text{R}^{16}$ ,  $\text{NR}^{16}\text{COR}^{16}$ ,  $\text{NR}^{16}\text{CO}_2\text{R}^{16}$ ,  $\text{CO}_2\text{R}^{16}$ ,  $\text{COR}^{16}$ ,  $\text{CONR}^{16}_2$ ,  $\text{S(O)}_2\text{R}^{16}$ ,  $\text{SONH}_2$ ,  $\text{S(O)R}^{16}$ ,  $\text{SO}_2\text{NR}^{16}\text{R}^{16}$ ,  $\text{NR}^{16}\text{S(O)}_2\text{R}^{16}$ , wherein the  $\text{C}_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group



consisting of  $-O-$ ,  $-N(R^{16})-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^{16}$  may be the same or different and is as defined below;

wherein  $R^{16}$  is hydrogen,  $C_{1-4}$  alkyl, or  $C_{1-4}$  haloalkyl;

5

wherein when X is  $NR^2$ ,  $R^2$  and  $R^3$  can form a 3 to 12 membered heterocyclyl ring, more preferably a 5, 6, 7, 8, 9, 10, 11 or 12 membered ring, wherein said ring can be partially saturated, unsaturated or fully saturated containing one to three heteroatoms; wherein the heterocyclyl group formed by  $R^2$  and  $R^3$  can  
 10 be optionally fused to one to three unsaturated, partially saturated or fully saturated 5 to 7 membered rings and contains from zero to three heteroatoms, any of said rings being optionally and independently substituted with one or more of  $C_{1-6}$  alkyl, halogen,  $C_{1-6}$  haloalkyl,  $OR^{22}$ ,  $SR^{22}$ ,  $NO_2$ ,  $CN$ ,  $NR^{22}R^{22}$ ,  $NR^{22}COR^{22}$ ,  $NR^{22}CONR^{22}R^{22}$ ,  $NR^{22}COR^{22}$ ,  $NR^{22}CO_2R^{22}$ ,  $CO_2R^{22}$ ,  $COR^{22}$ ,  
 15  $CONR^{22}_2$ ,  $S(O)_2R^{22}$ ,  $SONR^{22}_2$ ,  $S(O)R^{22}$ ,  $SO_2NR^{22}R^{22}$ ,  $NR^{22}S(O)_2R^{22}$ , wherein the  $C_{1-6}$  alkyl group optionally incorporates one or two insertions from  $-O-$ ,  $-N(R^{22})-$ ,  $-S(O)-$  and  $-S(O_2)-$  and wherein each  $R^{22}$  may be the same or different;

and wherein  $R^4$  is a six-membered carbocyclyl group or a five or six-membered  
 20 heterocyclyl group containing from 1 to 4 heteroatoms independently selected from N, S or O, wherein the optionally substituted six-membered carbocyclyl or five or six-membered heterocyclyl group is optionally fused to a partially saturated, unsaturated or fully saturated five to seven membered ring containing zero to three heteroatoms, and each substitutable carbon or hetero-atom in  $R^4$   
 25 including the optional fused ring, is optionally and independently substituted by one or more of halogen,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{1-12}$  haloalkyl,  $C_{3-12}$  carbocyclyl,  $C_{3-12}$  heterocyclyl,  $(CH_2)_nOR^{17}$ ,  $(CH_2)_nNR^{17}_2$ ,  $OR^{17}$ ,  $SR^{17}$ ,  $NO_2$ ,  $CN$ ,  $NR^{17}_2$ ,  $NR^{17}COR^{17}$ ,  $NR^{17}CONR^{17}_2$ ,  $NR^{17}COR^{17}$ ,  $NR^{17}CO_2R^{17}$ ,  $CO_2R^{17}$ ,  $COR^{17}$ ,  $CONR^{17}_2$ ,  $S(O)_2R^{17}$ ,  $SONR^{17}_2$ ,  $S(O)R^{17}$ ,  $SO_2NR^{17}_2$ , or

NR<sup>17</sup>S(O)<sub>2</sub>R<sup>17</sup>, wherein the C<sub>1-12</sub> alkyl group optionally contains one or more insertions selected from -O-, -N(R<sup>12</sup>)-, -S-, -S(O)- and -S(O<sub>2</sub>)-; and each saturated carbon in the optional fused ring is further optionally and independently substituted by =O, =S, NNR<sup>18</sup><sub>2</sub>, =N-OR<sup>18</sup>, =NNR<sup>18</sup>COR<sup>18</sup>,  
 5 =NNR<sup>18</sup>CO<sub>2</sub>R<sup>18</sup>, =NNSO<sub>2</sub>R<sup>18</sup>, or =NR<sup>18</sup>; and each substitutable nitrogen atom in R<sup>4</sup> is optionally substituted by R<sup>19</sup>, COR<sup>19</sup>, SO<sub>2</sub>R<sup>19</sup> or CO<sub>2</sub>R<sup>19</sup>; wherein n is 1 to 6, preferably n is 1, 2 or 3; preferably, wherein each substitutable carbon or hetero-atom in R<sup>4</sup> is optionally and independently substituted by one or more of C<sub>1-6</sub> alkyl, OR<sup>20</sup>, SR<sup>20</sup>, NO<sub>2</sub>, CN, NR<sup>20</sup><sub>2</sub>, NR<sup>20</sup>COR<sup>20</sup>, NR<sup>20</sup>CONR<sup>20</sup><sub>2</sub>,  
 10 NR<sup>20</sup>COR<sup>20</sup>, NHCO<sub>2</sub>R<sup>20</sup>, CO<sub>2</sub>R<sup>20</sup>, COR<sup>20</sup>, CONR<sup>20</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>20</sup>, SONR<sup>20</sup><sub>2</sub>, S(O)R<sup>20</sup>, SO<sub>2</sub>NR<sup>20</sup><sub>2</sub>, or NR<sup>20</sup>S(O)<sub>2</sub>R<sup>20</sup>;

wherein R<sup>20</sup> is hydrogen, C<sub>1-6</sub> alkyl, or C<sub>1-6</sub> haloalkyl;

15 wherein R<sup>17</sup> is hydrogen, C<sub>1-12</sub> alkyl, C<sub>3-12</sub> carbocyclyl or C<sub>3-12</sub> heterocyclyl, optionally substituted by one or more of C<sub>1-6</sub> alkyl, C<sub>3-12</sub> carbocyclyl, C<sub>3-12</sub> heterocyclyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>21</sup>, SR<sup>21</sup>, NO<sub>2</sub>, CN, NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CONR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CO<sub>2</sub>R<sup>21</sup>, CO<sub>2</sub>R<sup>21</sup>, COR<sup>21</sup>, CONR<sup>21</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>21</sup>, SONR<sup>21</sup><sub>2</sub>, S(O)R<sup>21</sup>, SO<sub>2</sub>NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>S(O)<sub>2</sub>R<sup>21</sup>, wherein  
 20 the C<sub>1-12</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -N(R<sup>21</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>21</sup> may be the same or different and is as defined below;

wherein two R<sup>17</sup> in NR<sup>17</sup><sub>2</sub> may optionally form a partially saturated, unsaturated  
 25 or fully saturated three to seven membered ring containing one to three heteroatoms, optionally and independently substituted by one or more of C<sub>1-6</sub> alkyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>21</sup>, SR<sup>21</sup>, NO<sub>2</sub>, CN, NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CONR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CO<sub>2</sub>R<sup>21</sup>, CO<sub>2</sub>R<sup>21</sup>, COR<sup>21</sup>, CONR<sup>21</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>21</sup>, SONR<sup>21</sup><sub>2</sub>, S(O)R<sup>21</sup>, SO<sub>2</sub>NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>S(O)<sub>2</sub>R<sup>21</sup>, wherein the C<sub>1-6</sub>

alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -N(R<sup>21</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>21</sup> may be the same or different and is as defined below;

- 5 wherein R<sup>18</sup> is hydrogen, C<sub>1-12</sub> alkyl, C<sub>3-12</sub> carbocyclyl or C<sub>3-12</sub> heterocyclyl, optionally substituted by one or more of C<sub>1-6</sub> alkyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>21</sup>, SR<sup>21</sup>, NO<sub>2</sub>, CN, NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CONR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CO<sub>2</sub>R<sup>21</sup>, CO<sub>2</sub>R<sup>21</sup>, COR<sup>21</sup>, CONR<sup>21</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>21</sup>, S(O)R<sup>21</sup>, SO<sub>2</sub>NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>S(O)<sub>2</sub>R<sup>21</sup>, wherein the C<sub>1-12</sub> alkyl group optionally incorporates one or two  
 10 insertions selected from the group consisting of -O-, -N(R<sup>21</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>21</sup> may be the same or different and is as defined below;

wherein R<sup>19</sup> is hydrogen, C<sub>6-12</sub> aryl, C<sub>1-6</sub> alkyl or C<sub>1-6</sub> haloalkyl;

wherein R<sup>21</sup> is hydrogen, C<sub>1-6</sub> alkyl, or C<sub>1-6</sub> haloalkyl;

15

and the pharmaceutically acceptable salts, and other pharmaceutically acceptable biohydrolyzable derivatives thereof, including esters, amides, carbamates, carbonates, ureides, solvates, hydrates, affinity reagents or prodrugs thereof.

20

For the avoidance of doubt when a group as defined above contains two or more radicals eg the radical R<sup>21</sup> as for example in the groups SO<sub>2</sub>NR<sup>21</sup>R<sup>21</sup> and NR<sup>3</sup>COR<sup>3</sup>, the two or more radicals i.e. R<sup>21</sup> may be the same or different.

- 25 For the purposes of this invention, alkyl relates to both straight chain and branched alkyl radicals of 1 to 12 carbon atoms, preferably 1 to 8 carbon atoms and most preferably 1 to 4 carbon atoms including but not limited to methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl n-pentyl, n-hexyl, n-heptyl, n-octyl. In particular, alkyl relates to a group having 1, 2, 3, 4,



5, 6, 7, 8, 9, 10, 11 or 12 carbon atoms. The term alkyl also encompasses cycloalkyl radicals including but not limited to cyclopropyl, cyclobutyl, CH<sub>2</sub>-cyclopropyl, CH<sub>2</sub>-cyclobutyl, cyclopentyl or cyclohexyl. In particular, cycloalkyl relates to a group having 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 carbon atoms. Cycloalkyl groups may be optionally substituted or fused to one or more carbocyclyl or heterocyclyl group. Haloalkyl relates to an alkyl radical as defined above preferably having 1 to 8 carbon atoms, preferably 1 to 4 carbon atoms substituted with one or more halide atoms for example one or more of F, Cl, Br or I, such as CH<sub>2</sub>CH<sub>2</sub>Br, CF<sub>3</sub> or CCl<sub>3</sub>.

10

The term "alkenyl" means a straight chain or branched alkylenyl radical of 2 to 12 carbon atoms, preferably 2 to 6 carbon atoms and most preferably 2 to 4 carbon atoms, and containing one or more carbon-carbon double bonds and includes but is not limited to ethylene, n-propyl-1-ene, n-propyl-2-ene, isopropylene, etc. In particular, alkenyl relates to a group having 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 carbon atoms. The term "alkynyl" means a straight chain or branched alkynyl radical of 2 to 12 carbon atoms, preferably 2 to 6 carbon atoms and most preferably 2 to 4 carbon atoms, and containing one or more carbon-carbon triple bonds and includes but is not limited to ethynyl, 2-methylethynyl etc. In particular, alkynyl relates to a group having 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 carbon atoms.

"Carbocyclyl" relates to a saturated, partly unsaturated or unsaturated 3-12 membered hydrocarbon ring preferably a 6-12 membered hydrocarbon ring, including cycloalkyl and aryl.

"Aryl" means an aromatic 3-12 membered hydrocarbon preferably a 6-12 membered hydrocarbon containing one ring or being fused to one or more

saturated or unsaturated rings including but not limited to phenyl, naphthyl, anthracenyl or phenanthracenyl.

“Heteroaryl” means an aromatic 3-12 membered aryl preferably a 6-12 membered aryl containing one or more heteroatoms selected from N, O or S and containing one ring or being fused to one or more saturated or unsaturated rings and;

“Heterocyclyl” means a 3-12 membered ring system preferably a 6-12 membered ring system containing one or more heteroatoms selected from N, O or S and includes heteroaryl. In particular the terms "carbocyclyl", "aryl", "heteroaryl" and "heterocyclyl" relate to a group having 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 carbon atoms.

The heterocyclyl system can contain one ring or may be fused to one or more saturated or unsaturated rings; the heterocyclyl can be fully saturated, partially saturated or unsaturated and includes but is not limited to heteroaryl and heterocarbocyclyl. Examples of carbocyclyl or heterocyclyl groups include but are not limited to cyclohexyl, phenyl, acridine, benzimidazole, benzofuran, benzothiophene, benzoxazole, benzothiazole, carbazole, cinnoline, dioxin, dioxane, dioxolane, dithiane, dithiazine, dithiazole, dithiolane, furan, imidazole, imidazoline, imidazolidine, indole, indoline, indolizine, indazole, isoindole, isoquinoline, isoxazole, isothiazole, morpholine, naphthyridine, oxazole, oxadiazole, oxathiazole, oxathiazolidine, oxazine, oxadiazine, phenazine, phenothiazine, phenoxazine, phthalazine, piperazine, piperidine, pteridine, purine, pyran, pyrazine, pyrazole, pyrazoline, pyrazolidine, pyridazine, pyridine, pyrimidine, pyrrole, pyrrolidine, pyrroline, quinoline, quinoxaline, quinazoline, quinolizine, tetrahydrofuran, tetrazine, tetrazole, thiophene,

thiadiazine, thiadiazole, thiatriazole, thiazine, thiazole, thiomorpholine, thianaphthalene, thiopyran, triazine, triazole, and trithiane.

For the purpose of the present invention, the term "fused" includes a polycyclic  
5 compound in which one ring contains one or more atoms preferably one, two or three atoms in common with one or more other ring.

Halogen means F, Cl, Br or I, preferably F.

10  $R^1$  is preferably a group of formula (II) or an optionally substituted five or six membered carbocyclyl or heterocyclyl group wherein the carbocyclyl or heterocyclyl group is optionally fused to one or more unsaturated rings.

When  $R^1$  is a substituted five or six membered carbocyclyl or heterocyclyl  
15 group it is preferably selected from optionally substituted phenyl, acridine, benzimidazole, benzofuran, benzothiophene, benzoxazole, benzothiazole, cyclohexyl furan, imidazole, indole, isoindole, isoquinoline, isoxazole, isothiazole, morpholine, naphthalene, oxazole, phenazine, phenothiazine, phenoxazine, piperazine, piperidine, pyrazole, pyridazine, pyridine, pyrrole,  
20 quinoline, quinolizine, tetrahydrofuran, tetrazine, tetrazole, thiophene, thiazole, thiomorpholine, thianaphthalene, thiopyran, triazine, triazole or trithiane.

When  $R^1$  is a group of formula (II), X is preferably a group  $NR^3$ , Y is preferably absent and one or more of  $R^2$  and  $R^3$  are preferably hydrogen alkyl  
25 or cycloalkyl, in particular, the group of formula (II) is preferably an alkylamino or cycloalkylamino group preferably selected from optionally substituted methylamino, ethylamino, propylamino, isopropylamino, butylamino, cyclobutylamino, pentylamino, cyclopentylamino, hexylamino, cyclohexylamino, heptylamino, cycloheptylamino, octylamino and

cyclooctylamino. In particular, X is an alkylamino or a cycloalkylamino group wherein the alkyl group has 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 carbon atoms and the cycloalkyl group has 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12 carbon atoms. R<sup>1</sup> may additionally be a group of formula (II) wherein X is NR<sup>3</sup> and R<sup>2</sup> and R<sup>3</sup> form a 5, 6, 7 or 8 membered ring, said ring being partially, saturated, fully saturated or unsaturated and optionally substituted as previously discussed.

As discussed above, R<sup>1</sup> can be optionally substituted at any position on the alkylamino, cycloalkyl amino, carbocyclyl, heterocyclyl or optional fused ring.

10

R<sup>1</sup> is preferably substituted with one or more of OR<sup>24</sup>, halogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>1-6</sub> haloalkyl, C<sub>1-6</sub> alkylaryl, C<sub>1-6</sub> alkylheterocyclyl, (CH<sub>2</sub>)<sub>n</sub>OR<sup>24</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>24</sup><sub>2</sub>, SR<sup>24</sup>, NO<sub>2</sub>, CN, NR<sup>24</sup><sub>2</sub>, CO<sub>2</sub>R<sup>24</sup>, NR<sup>24</sup>C(O)R<sup>24</sup>, NR<sup>24</sup>S(O)<sub>2</sub>R<sup>24</sup>, COR<sup>24</sup>, CONR<sup>24</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>24</sup>, S(O)R<sup>24</sup> or SO<sub>2</sub>NR<sup>24</sup><sub>2</sub>;

15 wherein R<sup>24</sup> is hydrogen, C<sub>1-4</sub> alkyl or C<sub>6-12</sub> aryl preferably phenyl, or C<sub>5-12</sub> heterocyclyl preferably pyridine, and n is 1, 2, 3, 4, 5 or 6.

wherein two R<sup>24</sup> in NR<sup>24</sup><sub>2</sub> may optionally form a partially saturated, unsaturated or fully saturated three to seven membered ring containing one to three heteroatoms, said ring is preferably independently substituted with one or more of halogen, C<sub>1-12</sub> alkyl, C<sub>2-12</sub> alkenyl, C<sub>2-12</sub> alkynyl, C<sub>1-12</sub> haloalkyl, C<sub>3-12</sub> carbocyclyl, C<sub>3-12</sub> heterocyclyl, OR<sup>25</sup>, SR<sup>25</sup>, NO<sub>2</sub>, CN, NR<sup>25</sup><sub>2</sub>, NR<sup>25</sup>COR<sup>25</sup>, NR<sup>25</sup>CONR<sup>25</sup><sub>2</sub>, NR<sup>25</sup>COR<sup>25</sup>, NR<sup>25</sup>CO<sub>2</sub>R<sup>25</sup>, CO<sub>2</sub>R<sup>25</sup>, COR<sup>25</sup>, CONR<sup>25</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>25</sup>, SONR<sup>25</sup><sub>2</sub>, S(O)R<sup>25</sup>, SO<sub>2</sub>NR<sup>25</sup><sub>2</sub>, or NR<sup>25</sup>S(O)<sub>2</sub>R<sup>25</sup>; and each saturated carbon in the optional ring is further optionally and independently substituted by =O, =S, NNR<sup>26</sup><sub>2</sub>, =N-OR<sup>26</sup>, =NNR<sup>26</sup>COR<sup>26</sup>, =NNR<sup>26</sup>CO<sub>2</sub>R<sup>26</sup>, =NNSO<sub>2</sub>R<sup>26</sup>, or =NR<sup>26</sup>; and each substitutable nitrogen atom is optionally substituted by R<sup>27</sup>, COR<sup>27</sup>, SO<sub>2</sub>R<sup>27</sup> or CO<sub>2</sub>R<sup>27</sup>;

wherein  $R^{25}$  is hydrogen,  $C_{1-12}$  alkyl,  $C_{6-12}$  carbocyclyl or  $C_{5-12}$  heterocyclyl, optionally substituted by one or more of  $C_{1-6}$  alkyl, halogen,  $C_{1-6}$  haloalkyl,  $OR^{28}$ ,  $SR^{28}$ ,  $NO_2$ ,  $CN$ ,  $NR^{28}R^{28}$ ,  $NR^{28}COR^{28}$ ,  $NR^{28}CONR^{28}R^{28}$ ,  $NR^{28}COR^{28}$ ,  $NR^{28}CO_2R^{28}$ ,  $CO_2R^{28}$ ,  $COR^{28}$ ,  $CONR^{28}_2$ ,  $S(O)_2R^{28}$ ,  $\dot{S}ONR^{28}_2$ ,  $S(O)R^{28}$ ,  
 5  $SO_2NR^{28}R^{28}$ ,  $NR^{28}S(O)_2R^{28}$ , wherein the  $C_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-O-$ ,  $-N(R^{28})-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^{28}$  may be the same or different and is as defined below;

10 wherein  $R^{26}$  is hydrogen,  $C_{1-12}$  alkyl,  $C_{6-12}$  carbocyclyl or  $C_{5-12}$  heterocyclyl, optionally substituted by one or more of  $C_{1-6}$  alkyl, halogen,  $C_{1-6}$  haloalkyl,  $OR^{28}$ ,  $SR^{28}$ ,  $NO_2$ ,  $CN$ ,  $NR^{28}R^{28}$ ,  $NR^{28}COR^{28}$ ,  $NR^{28}CONR^{28}R^{28}$ ,  $NR^{28}COR^{28}$ ,  $NR^{28}CO_2R^{28}$ ,  $CO_2R^{28}$ ,  $COR^{28}$ ,  $CONR^{28}_2$ ,  $S(O)_2R^{28}$ ,  $S(O)R^{28}$ ,  $SO_2NR^{28}R^{28}$ ,  $NR^{28}S(O)_2R^{28}$ , wherein the  $C_{1-12}$  alkyl group optionally incorporates one or two  
 15 insertions selected from the group consisting of  $-O-$ ,  $-N(R^{28})-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^{28}$  may be the same or different and is as defined below;

wherein  $R^{27}$  is hydrogen,  $C_{1-6}$  alkyl,  $C_{1-6}$  haloalkyl or  $C_{6-12}$  aryl;

20 wherein  $R^{28}$  is hydrogen,  $C_{1-6}$  alkyl, or  $C_{1-6}$  haloalkyl.

$R^4$  is preferably selected from phenyl, cyclohexyl, acridine, benzimidazole, benzofuran, benzothiophene, benzoxazole, benzothiazole, indole, isoindole, indolizine, indazole, isoindole, isoquinoline, morpholine, naphthalene,  
 25 phenazine, phenothiazine, phenoxazine, piperazine, piperidine, pyridazine, pyridine, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, quinoline, quinolizine, tetrazine, thiomorpholine, thianaphthalene, thiopyran, triazine, trithiane, furan, imidazole, isoxazole, isothiazole, oxazole, oxadiazole, oxathiazole, pyrazole, pyrrole, tetrazole, thiophene, thiadiazole, thiatriazole, thiazole or triazole.



As discussed above,  $R^4$  can be optionally substituted at any position on the carbocyclyl, heterocyclyl or optional fused ring. Preferably, each substitutable carbon or hetero-atom in  $R^4$  is optionally and independently substituted by one or more of  $C_{1-6}$  alkyl,  $OR^{20}$ ,  $SR^{20}$ ,  $NO_2$ ,  $CN$ ,  $NR^{20}_2$ ,  $NR^{20}COR^{20}$ ,  $NR^{20}CONR^{20}_2$ ,  $NR^{20}COR^{20}$ ,  $NHCO_2R^{20}$ ,  $CO_2R^{20}$ ,  $COR^{20}$ ,  $CONR^{20}_2$ ,  $S(O)_2R^{20}$ ,  $SONR^{20}_2$ ,  $S(O)R^{20}$ ,  $SO_2NR^{20}_2$ , or  $NR^{20}S(O)_2R^{20}$ ;

wherein  $R^{20}$  is hydrogen,  $C_{1-6}$  alkyl, or  $C_{1-6}$  haloalkyl.

10

When  $R^4$  is a six-membered carbocyclyl or heterocyclyl group,  $R^4$  is preferably substituted with one or more of  $OR^{29}$ ,  $NR^{29}_2$ ,  $SR^{29}$ ,  $(CH_2)_nOR^{29}$ ,  $(CH_2)_nNR^{29}_2$ , halogen,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl, haloalkyl,  $NO_2$ ,  $CN$ ,  $NR^{29}C(O)R^{29}$ ,  $NR^{29}S(O)_2R^{29}$ ,  $CO_2R^{29}$ ,  $COR^{29}$ ,  $CONR^{29}_2$ ,  $S(O)_2R^{29}$ ,  $S(O)R^{29}$  or  $SO_2NR^{29}_2$ ;

15

wherein  $R^{29}$  is hydrogen,  $C_{1-4}$  alkyl,  $C_{5-12}$  heterocyclyl or  $C_{6-12}$  aryl preferably phenyl, and  $n$  is 1, 2, 3, 4, 5 or 6.

wherein two  $R^{29}$  in  $NR^{29}_2$  may optionally form a partially saturated, unsaturated or fully saturated five to seven membered ring containing one to three heteroatoms, optionally and independently substituted with one or more of halogen,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{1-12}$  haloalkyl,  $C_{6-12}$  carbocyclyl,  $C_{5-12}$  heterocyclyl,  $OR^{30}$ ,  $SR^{30}$ ,  $NO_2$ ,  $CN$ ,  $NR^{30}_2$ ,  $NR^{30}COR^{30}$ ,  $NR^{30}CONR^{30}_2$ ,  $NR^{30}COR^{30}$ ,  $NR^{30}CO_2R^{30}$ ,  $CO_2R^{30}$ ,  $COR^{30}$ ,  $CONR^{30}_2$ ,  $S(O)_2R^{30}$ ,  $SONR^{30}_2$ ,  $S(O)R^{30}$ ,  $SO_2NR^{30}_2$ , or  $NR^{30}S(O)_2R^{30}$ ; and each saturated carbon in the optional ring is further optionally and independently substituted by  $=O$ ,  $=S$ ,  $NNR^{31}_2$ ,  $=N-OR^{31}$ ,  $=NNR^{31}COR^{31}$ ,  $=NNR^{31}CO_2R^{31}$ ,  $=NNSO_2R^{31}$ , or  $=NR^{31}$ ; and each substitutable nitrogen atom is optionally substituted by  $R^{32}$ ,  $COR^{32}$ ,  $SO_2R^{32}$  or  $CO_2R^{32}$ ;

25


wherein  $R^{30}$  is hydrogen,  $C_{1-12}$  alkyl,  $C_{6-12}$  carbocyclyl or  $C_{5-12}$  heterocyclyl, optionally substituted by one or more of  $C_{1-6}$  alkyl, halogen,  $C_{1-6}$  haloalkyl,  $OR^{33}$ ,  $SR^{33}$ ,  $NO_2$ ,  $CN$ ,  $NR^{33}R^{33}$ ,  $NR^{33}COR^{33}$ ,  $NR^{33}CONR^{33}R^{33}$ ,  $NR^{33}COR^{33}$ ,  $NR^{33}CO_2R^{33}$ ,  $CO_2R^{33}$ ,  $COR^{33}$ ,  $CONR^{33}_2$ ,  $S(O)_2R^{33}$ ,  $SONR^{33}_2$ ,  $S(O)R^{33}$ ,  $SO_2NR^{33}R^{33}$ ,  $NR^{33}S(O)_2R^{33}$ , wherein the  $C_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-O-$ ,  $-N(R^{33})-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^{33}$  may be the same or different and is as defined below;

10 wherein  $R^{31}$  is hydrogen,  $C_{1-12}$  alkyl,  $C_{6-12}$  carbocyclyl or  $C_{5-12}$  heterocyclyl, optionally substituted by one or more of  $C_{1-6}$  alkyl, halogen,  $C_{1-6}$  haloalkyl,  $OR^{33}$ ,  $SR^{33}$ ,  $NO_2$ ,  $CN$ ,  $NR^{33}R^{33}$ ,  $NR^{33}COR^{33}$ ,  $NR^{33}CONR^{33}R^{33}$ ,  $NR^{33}COR^{33}$ ,  $NR^{33}CO_2R^{33}$ ,  $CO_2R^{33}$ ,  $COR^{33}$ ,  $CONR^{33}_2$ ,  $S(O)_2R^{33}$ ,  $S(O)R^{33}$ ,  $SO_2NR^{33}R^{33}$ ,  $NR^{33}S(O)_2R^{33}$ , wherein the  $C_{1-12}$  alkyl group optionally incorporates one or two  
15 insertions selected from the group consisting of  $-O-$ ,  $-N(R^{33})-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^{21}$  may be the same or different and is as defined below;

wherein  $R^{32}$  is hydrogen,  $C_{6-12}$  aryl,  $C_{1-6}$  alkyl or  $C_{1-6}$  haloalkyl;

20 wherein  $R^{33}$  is hydrogen,  $C_{1-6}$  alkyl, or  $C_{1-6}$  haloalkyl.

When  $R^4$  is a five-membered heterocyclyl, it is preferably a group





Wherein A, X<sup>2</sup>, Y<sup>2</sup> or Z are independently selected from N, O, C, S and M is C or N, wherein one, two, three or four of A, X<sup>2</sup>, Y<sup>2</sup>, Z and M is other than C, preferably R<sup>4</sup> is furan, imidazole, isoxazole, isothiazole, oxazole, oxadiazole, oxatriazole, pyrazole, pyrrole, tetrazole, thiophene, thiadiazole, thiatriazole, 5 thiazole or triazole;

R<sup>34</sup>, R<sup>35</sup>, R<sup>36</sup> or R<sup>37</sup> are independently selected from a lone electron pair, hydrogen, halogen, C<sub>1-12</sub> alkyl, C<sub>1-12</sub> haloalkyl, OR<sup>38</sup>, SR<sup>38</sup>, NO<sub>2</sub>, CN, NR<sup>38</sup><sub>2</sub>, NR<sup>38</sup>COR<sup>38</sup>, NR<sup>38</sup>CONR<sup>38</sup><sub>2</sub>, NR<sup>38</sup>COR<sup>38</sup>, NR<sup>38</sup>CO<sub>2</sub>R<sup>38</sup>, (CH<sub>2</sub>)<sub>n</sub>OR<sup>38</sup>, 10 (CH<sub>2</sub>)<sub>n</sub>NR<sup>38</sup><sub>2</sub>, CO<sub>2</sub>R<sup>38</sup>, COR<sup>38</sup>, CONR<sup>38</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>38</sup>, SONR<sup>38</sup><sub>2</sub>, S(O)R<sup>38</sup>, SO<sub>2</sub>NR<sup>38</sup><sub>2</sub>, or NHS(O)<sub>2</sub>R<sup>38</sup>;

wherein n is 1 to 6, preferably n is 1, 2 or 3;

or wherein any two of R<sup>34</sup>, R<sup>35</sup>, R<sup>36</sup> or R<sup>37</sup> may optionally form a partially 15 saturated, unsaturated or fully saturated five to seven membered ring containing zero to three heteroatoms, each saturated carbon in the optional fused ring is further optionally and independently substituted with one or more of halogen, C<sub>1-12</sub> alkyl, C<sub>2-12</sub> alkenyl, C<sub>2-12</sub> alkynyl, C<sub>1-12</sub> haloalkyl, C<sub>6-12</sub> carbocyclyl, C<sub>5-12</sub> heterocyclyl, OR<sup>38</sup>, SR<sup>38</sup>, NO<sub>2</sub>, CN, NR<sup>38</sup><sub>2</sub>, NR<sup>38</sup>CONR<sup>38</sup><sub>2</sub>, NR<sup>38</sup>COR<sup>38</sup>, 20 NR<sup>38</sup>CO<sub>2</sub>R<sup>38</sup>, (CH<sub>2</sub>)<sub>n</sub>OR<sup>38</sup>, (CH<sub>2</sub>)<sub>n</sub>NR<sup>38</sup><sub>2</sub>, CO<sub>2</sub>R<sup>38</sup>, COR<sup>38</sup>, CONR<sup>38</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>38</sup>, SONR<sup>38</sup><sub>2</sub>, S(O)R<sup>38</sup>, SO<sub>2</sub>NR<sup>38</sup><sub>2</sub>, or NR<sup>38</sup>S(O)<sub>2</sub>R<sup>38</sup>; and each saturated carbon in the optional fused ring is further optionally and independently substituted by =O, =S, NNR<sup>39</sup><sub>2</sub>, =N-OR<sup>39</sup>, =NNR<sup>39</sup>COR<sup>39</sup>, =NNR<sup>39</sup>CO<sub>2</sub>R<sup>39</sup>, =NNSO<sub>2</sub>R<sup>39</sup>, or =NR<sup>39</sup>; and each substitutable nitrogen atom in R<sup>4</sup> is optionally substituted by 25 R<sup>40</sup>, COR<sup>40</sup>, SO<sub>2</sub>R<sup>40</sup> or CO<sub>2</sub>R<sup>40</sup>;

wherein n is 1 to 6, preferably n is 1, 2 or 3;

wherein R<sup>38</sup> is hydrogen, C<sub>1-12</sub> alkyl, C<sub>6-12</sub> carbocyclyl or C<sub>5-12</sub> heterocyclyl, optionally substituted by one or more of C<sub>1-6</sub> alkyl, halogen, C<sub>1-6</sub> haloalkyl,

OR<sup>41</sup>, SR<sup>41</sup>, NO<sub>2</sub>, CN, NR<sup>41</sup>R<sup>41</sup>, NR<sup>41</sup>CONR<sup>41</sup>R<sup>41</sup>, NR<sup>41</sup>COR<sup>41</sup>, NR<sup>41</sup>CO<sub>2</sub>R<sup>41</sup>,  
 CO<sub>2</sub>R<sup>41</sup>, COR<sup>41</sup>, CONR<sup>41</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>41</sup>, SONR<sup>41</sup><sub>2</sub>, S(O)R<sup>41</sup>, SO<sub>2</sub>NR<sup>41</sup>R<sup>41</sup>,  
 NR<sup>41</sup>S(O)<sub>2</sub>R<sup>41</sup>, wherein the C<sub>1-12</sub> alkyl group optionally incorporates one or two  
 5 S(O)<sub>2</sub>-, wherein each R<sup>41</sup> may be the same or different and is as defined below;


























wherein R<sup>39</sup> is hydrogen, C<sub>1-12</sub> alkyl, carbocyclyl or heterocyclyl, optionally  
 substituted by one or more of C<sub>1-6</sub> alkyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>41</sup>, SR<sup>41</sup>,  
 NO<sub>2</sub>, CN, NR<sup>41</sup>R<sup>41</sup>, NR<sup>41</sup>COR<sup>41</sup>, NR<sup>41</sup>CONR<sup>41</sup>R<sup>41</sup>, NR<sup>41</sup>COR<sup>41</sup>, NR<sup>41</sup>CO<sub>2</sub>R<sup>41</sup>,  
 10 CO<sub>2</sub>R<sup>41</sup>, COR<sup>41</sup>, CONR<sup>41</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>41</sup>, S(O)R<sup>41</sup>, SO<sub>2</sub>NR<sup>41</sup>R<sup>41</sup>, NR<sup>41</sup>S(O)<sub>2</sub>R<sup>41</sup>,  
 wherein the C<sub>1-12</sub> alkyl group optionally incorporates one or two insertions  
 selected from the group consisting of -O-, -N(R<sup>41</sup>)-, -S(O)- and -S(O)<sub>2</sub>-,  
 wherein each R<sup>41</sup> may be the same or different and is as defined below;


























15 wherein R<sup>40</sup> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl or C<sub>6-12</sub> aryl.


wherein R<sup>41</sup> is hydrogen, C<sub>1-6</sub> alkyl, or C<sub>1-6</sub> haloalkyl.

More preferably R<sup>34</sup>, R<sup>35</sup>, R<sup>36</sup> or R<sup>37</sup> are independently selected from a lone  
 20 electron pair, hydrogen, halogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, OR<sup>42</sup>, SR<sup>42</sup>, CN,  
 NR<sup>42</sup><sub>2</sub>, NR<sup>42</sup>COR<sup>42</sup>, CO<sub>2</sub>R<sup>42</sup>, COR<sup>42</sup>, CONR<sup>42</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>42</sup>, or S(O)R<sup>42</sup>;  
 wherein R<sup>42</sup> is hydrogen, C<sub>1-4</sub> alkyl, preferably methyl or ethyl or carbocyclyl,  
 preferably phenyl.

25 Representative compounds according to the first aspect of the invention are  
 illustrated below.







The compounds of the first aspect may be provided as a salt, preferably as a pharmaceutically acceptable salt of compounds of formula (I). Examples of pharmaceutically acceptable salts of these compounds include those derived from organic acids such as acetic acid, malic acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid, mandelic acid, methanesulphonic acid, benzenesulphonic acid and p-toluenesulphonic acid, mineral acids such as hydrochloric and sulphuric acid and the like, giving methanesulphonate, benzenesulphonate, p-toluenesulphonate, hydrochloride and sulphate, and the like, respectively or those derived from bases such as organic and inorganic bases. Examples of suitable inorganic bases for the formation of salts of compounds for this invention include the hydroxides, carbonates, and bicarbonates of ammonia, lithium, sodium, calcium, potassium, aluminium, iron, magnesium, zinc and the like. Salts can also be formed with suitable organic bases. Such bases suitable for the formation of pharmaceutically acceptable base addition salts with compounds of the present invention include organic bases, which are nontoxic and strong enough to form salts. Such organic bases are already well known in the art and may include amino acids such as arginine and lysine, mono-, di-, or trihydroxyalkylamines such as

mono-, di-, and triethanolamine, choline, mono-, di-, and trialkylamines, such as methylamine, dimethylamine, and trimethylamine, guanidine; N-methylglucosamine; N-methylpiperazine; morpholine; ethylenediamine; N-benzylphenethylamine; tris(hydroxymethyl) aminomethane; and the like.

5

Salts may be prepared in a conventional manner using methods well known in the art. Acid addition salts of said basic compounds may be prepared by dissolving the free base compounds according to the first aspect of the invention in aqueous or aqueous alcohol solution or other suitable solvents  
10 containing the required acid. Where a compound of the invention contains an acidic function, a base salt of said compound may be prepared by reacting said compound with a suitable base. The acid or base salt may separate directly or can be obtained by concentrating the solution e.g. by evaporation. The compounds of this invention may also exist in solvated or hydrated forms.

15


The invention also extends to a prodrug of the aforementioned compounds such as an ester or amide thereof. A prodrug is any compound that may be converted under physiological conditions or by solvolysis to any of the compounds of the invention or to a pharmaceutically acceptable salt of the compounds of the  
20 invention. A prodrug may be inactive when administered to a subject but is converted *in vivo* to an active compound of the invention.

The compounds of the invention may contain one or more asymmetric carbon atoms and may exist in racemic and optically active forms. The compounds of  
25 the invention may exist in trans or cis form. The first aspect of the invention covers all of these compounds.

The second aspect of the invention provides a process for the manufacture of a compound of formula (I) wherein  $R^1$  is a group of formula (II) as defined in the



first aspect of the invention comprising the condensation of an intermediate (III) with an intermediate (IV).




- 5 wherein  $\text{R}^2$  and  $\text{R}^4$  are as defined in the first aspect of the invention;  $\text{L}^1$  and  $\text{L}^2$  are independently a leaving group wherein  $\text{L}^1$  and  $\text{L}^2$  together form a condensation product.

According to the process, a compound of the general formula (III), undergoes a  
 10 condensation reaction with the compound of the general formula (IV), to form a compound of general formula I. In formulae (III) and (IV),  $\text{L}^1$  and  $\text{L}^2$  represent radicals that together form a condensation product, e.g. H and OH or H and Cl. Preferably  $\text{L}^1$  is OH,  $\text{OR}^{50}$ , OM, Cl, Br or I wherein  $\text{R}^{50}$  is  $\text{C}_{1-6}$  alkyl, preferably methyl or ethyl and M is a metal, preferably Na, Li, K, Ca, Mg or Ba, and  $\text{L}^2$  is  
 15 preferably hydrogen or M. The condensation reaction occurs in a solution, preferably in a polar aprotic solvent such as e.g. dimethylformamide or dichloromethane. The condensation reaction may occur under the influence of coupling agents such as, for instance WSC·HCl, DCC, benzotriazole-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate (BOP), PyBrOP, etc.,  
 20 and in the presence of a tertiary amine (e.g. triethylamine) and 1-hydroxybenzotriazole (HOBT). Alternatively, the acid (III) may be first converted to an acid chloride by treatment with, for example, oxalyl chloride or thionyl chloride, and then without purification, reacted with, e.g. amines of formula (IV).



The third aspect of the invention provides a compound of formula (III)



(III)

wherein  $R^4$  is as defined in the first aspect of the invention and


$L^1$  is OH,  $OR^{50}$ , OM, Cl, Br, I;

5  $R^{50}$  is  $C_{1-6}$  alkyl, preferably methyl or ethyl; and

M is metal, preferably Na, Li, K, Ca, Mg, Ba.

A compound of formula (III) may undergo one or more further reactions to provide a different compound of formula (III). For example, a compound may  
10 undergo a hydrolysis, reduction, oxidation, elimination, substitution and/or addition reaction.

The fourth aspect of the invention provides a process for the manufacture of a compound of formula (V) comprising removal of group  $R^{51}$  from an  
15 intermediate (VI)



wherein  $L^3$  is  $R^1$  or  $L^1$ ,

wherein  $R^1$  is as defined in the first aspect and

$L^1$  is as defined in the second or third aspect;

20  $R^4$  is as defined in the first aspect, and

$R^{51}$  is an amino protecting group. The pyrrole nitrogen can be protected using any protection known in the art.  $R^{51}$  can therefore include  $R^{52}SO_2$ ,  $R^{52}C(O)$ ,

- $R^{52}_3\text{Si}$ ,  $R^{52}\text{OCH}_2$ ,  $(R^{52})_2\text{NSO}_2$ ,  $(R^{52})_2\text{NC(O)}$ ,  $R^{52}\text{OC(O)}$ ,  $R^{52}(R^{52}\text{O})\text{CH}$ ,  $R^{52}\text{CH}_2\text{CH}_2$ ,  $R^{52}\text{CH}_2$ ,  $\text{PhC(O)CH}_2$ ,  $\text{CH}_2=\text{CH}$ ,  $\text{ClCH}_2\text{CH}_2$ ,  $\text{Ph}_3\text{C}$ ,  $\text{Ph}_2(4\text{-pyridyl})\text{C}$ ,  $\text{Me}_2\text{N}$ ,  $\text{HO-CH}_2$ ,  $R^{52}\text{OCH}_2$ ,  $(R^{52})_3\text{SiOCH}_2$ ,  $(R^{52}\text{O})_2\text{CH}$ ,  $t\text{-BuOC(O)CH}_2$ ,  $\text{Me}_2\text{NCH}_2$ , and tetrahydropyranylamine,
- 5 wherein  $R^{52}$  is  $\text{C}_{1-6}$  alkyl or  $\text{C}_{6-12}$  aryl.

More preferably  $R^{51}$  is sulfonamide, most preferably benzenesulfonamide,  $(R^{52})_2\text{NSO}_2$ , and  $(R^{52})_2\text{NC(O)}$ ,


- 10 Removal of the protecting group can be afforded using conditions relevant to the protecting group used i.e. sulfonamide or amide protection can be removed by hydrolysis under basic conditions for example sodium hydroxide in water-ethanol, and silyl protection can be removed under acidic conditions for example TFA, HCl or using a source of fluoride, for example TBAF.

15

It will be appreciated that when  $L^3$  is  $R^1$  the deprotection will afford directly the compound of formula (I).

- It will be further appreciated than when  $L^1$  is  $\text{OR}^{50}$  the hydrolytic removal of
- 20  $R^{51}$  under basic conditions is accompanied by hydrolysis of the ester functionality to afford compound (III) where  $L^1$  is OM, or OH after acidification of the reaction mixture.

The fifth aspect of the invention provides a compound of formula (VI)



(VI)


25

wherein  $R^4$  is as defined in the first aspect,

$L^3$  is as defined in the fourth aspect; and

$R^{51}$  is an amino protecting group as defined in the fourth aspect.

The sixth aspect of the invention provides a process for the manufacture of a  
 5 compound of formula (VI) as defined in the fifth aspect of the invention  
 comprising a a) reaction of a compound of formula (VII) with stannane  $R^4$ -  
 $Sn(R^{53})_3$  in the presence of a palladium catalyst or b) reaction of a compound of  
 formula (VII) with boronic acid or ester  $R^4$ - $B(OR^{54})_2$  in a presence of a suitable  
 palladium catalyst or c) reaction of a compound of formula (VII) with silane  
 10  $R^4$ - $Si(R^{55})_3$  in the presence of a palladium catalyst;



wherein  $R^4$  is as defined in the first aspect,

$L^3$  is as defined in the fourth aspect,

$R^{51}$  is an amino protecting group defined in the fourth aspect.

15  $X^3$  is F, Cl, Br I or  $CF_3SO_3$  preferably I or Br,

and wherein  $R^{53}$  is independently  $C_{1-6}$  alkyl;

$R^{54}$  is independently hydrogen or  $C_{1-6}$  alkyl or wherein two  $R^{54}$  groups together  
 optionally form a five, six or seven membered ring with the boron and oxygen  
 20 atoms, wherein the ring is optionally substituted with one or more  $C_{1-6}$  alkyl  
 group. Preferably,  $R^{54}$  is hydrogen or both  $R^{54}$  groups form the group -  
 $C(CH_3)_2-C(CH_3)_2$ ;

and  $R^{55}$  is independently  $C_{1-6}$  alkyl, F, OH.


Suitable catalysts for the purpose of this invention include  $(PPh_3)_2PdCl_2$ ,  $(PPh_3)_4Pd$ ,  $Pd(OAc)_2$ ,  $[PdCl(\eta^3-C_3H_5)]_2$ ,  $Pd_2(dba)_3$ ,  $Pd(dba)_2$  (dba = dibenzylidenacetone) and/or  $Pd/P(t-Bu)_3$ .

- 5 It will be appreciated that the reaction set out as option a) for the sixth aspect is a Stille reaction, which can be carried out according to Stille *Angew. Chem., Int.ed, Engl.* 1986, 25, 508; Mitchell *Synthesis*, 1992, 803, or Littke et al. *J. Am. Chem. Soc.* 2002, 124, 6343.
- 10 The reaction set out as option b) for the sixth aspect is a Suzuki reaction which can be carried out according to Suzuki *Pure Appl. Chem.* 1991, 63, 419 or Littke *J. Am. Chem. Soc.* 2000, 122, 4020

- It will be appreciated that the reaction set out as option c) for the sixth aspect is
- 15 a Hiyama reaction which can be carried out according to Hatanaka et al. *J. Org. Chem.* 1988, 53, 918, Hatanaka et al. *Synlett*, 1991, 845, Tamao et al. *Tetrahedron Lett.* 1989, 30, 6051 or Denmark et al. *Org. Lett.* 2000, 2, 565, *ibid.* 2491.

- 20 It will further be appreciated than when  $R^{51}$  is replaced with hydrogen the process of the sixth aspect yields a compound of formula (III) as defined in the fourth aspect of the invention.

The seventh aspect of the invention provides a compound of formula (VII)




(VII)

25

wherein  $L^3$  is as defined in the fourth aspect;

$R^{51}$  is an amino protecting group as defined in the fourth aspect;  
and  $X^3$  is as defined in the sixth aspect.

The eighth aspect of the invention provides a process for the manufacture of a  
5 compound of formula (VII) comprising protection of the pyrrole nitrogen of a  
compound of formula (VIII).




wherein  $L^3$  is as defined in the fourth aspect;  
10  $R^{51}$  is an amino protecting group defined in the fourth aspect, and  
 $X^3$  is as defined in the sixth aspect

Conditions for the introduction of the protecting group  $R^{51}$  will depend upon  
the protecting group used. Compound (VII) can be produced by the initial  
15 formation of the relevant salt, for example by treatment with BuLi in THF or  
NaH in DMF, followed by reaction of the salt with an electrophile such as  
sulfonyl halide, or acid chloride. Alternatively a compound of formula (VII)  
can be produced by the direct reaction of compound (VIII) with an electrophile  
such as benzenesulfonyl halide, preferably benzenesulfonyl chloride. This  
20 reaction is preferably carried out in the presence of base (such as sodium  
hydroxide) and a phase transfer catalyst such as tetra-n-butylammonium  
bromide or tetra-n-butylammonium hydrogen sulphate.

The ninth aspect of the invention provides a compound of formula (VIII)






(VIII)

wherein  $L^3$  is as defined in the fourth aspect and  $X^3$  is as defined in the sixth aspect.


- 5 The tenth aspect of the invention provides a process for the production of a compound of formula (VIII) by the introduction of an  $X^3$  group into a compound of formula (IX). Compound (VIII) can be produced from compound (IX) by halogenation under anhydrous conditions or by reaction with ICl under basic conditions (such as pyridine or  $i\text{-Pr}_2\text{NEt}$  in a chlorinated solvent such as
- 10  $\text{CH}_2\text{Cl}_2$ ,  $\text{CHCl}_3$ ,  $\text{CCl}_4$ ) or NBS in an anhydrous solvent such as  $\text{CH}_2\text{Cl}_2$ ,  $\text{CHCl}_3$ ,  $\text{CCl}_4$ ). Where  $X^3$  is iodine, it may preferably be introduced by direct action of  $\text{I}_2$  on (IX) in the presence of a strong base such as sodium hydroxide or potassium hydroxide in anhydrous solvent such as dimethylformamide.



- 15 wherein  $L^3$  is as defined in the fourth aspect and  $X^3$  is as defined in the sixth aspect.

Preparation of compound of general formula (IX) wherein  $L^3$  is carbocyclyl or heterocyclyl has been disclosed in GB0305142.2.

The eleventh aspect of the invention provides compound of formula (IX)




(IX)

wherein  $L^3$  is as defined in the fourth aspect with the exception that  $L^3$  is not carbocyclyl or heterocyclyl;

5

in particular wherein  $L^3$  is a group  $L^1$  as defined in the second aspect or a group  $R^1$ , wherein  $R^1$  is a group of formula (II)




(II)

wherein  $X$  is  $NR^3$ ,  $O$ ,  $S$  or  $(CR^{22}R^{22})_n$ ,  $Y$  is absent or is  $NR^{23}$ ,  $O$  or  $(CR^{23}R^{23})_n$ ,

10  $R^2$  is optionally substituted  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{3-12}$  carbocyclyl or  $C_{3-12}$  heterocyclyl as defined in the first aspect.

The twelfth aspect of the invention provides an alternative process for the production of a compound of formula (VII) by the introduction of the  $X^3$  group

15 to a compound of formula (X).



(IX)

(X)

(VII)

wherein  $R^{51}$  and  $L^3$  are as defined in the fourth aspect and  $X^3$  is as defined in the sixth aspect.

20 In particular where  $R^{51}$  is a silyl group, introduction of  $R^{51}$  occurs prior to the introduction of  $X^3$ . Preparation of compound of general formula (X) wherein  $L^3$  is carbocyclyl or heterocyclyl has been disclosed in GB0305142.2


Thus, a skilled person will appreciate that the actual synthetic sequence to prepare compound (VII) will depend on the type of protecting group  $R^{51}$  used, i.e. the compound (VII) can be prepared by the process of the twelfth aspect or  
 5 by the processes set out in the eighth and tenth aspects.

The thirteenth aspect of the invention provides a compound of formula (X)



wherein  $L^3$  is as defined in the eleventh aspect and  
 10 wherein  $R^{51}$  is an amino protecting group as defined in the fourth aspect.

The fourteenth aspect of the invention provides a method for preparation of compound of formula (IX) by the acid-catalysed hydrolysis of nitrile (XI) in the presence of alcohol, preferably methanol or ethanol.




15

wherein  $L^3$  is  $OR^{50}$ ;  
 and  $R^{50}$  is as defined in the third aspect of the invention. Preferably  $R^{50}$  is methyl or ethyl.

20 The acid-catalysed hydrolysis of compound (IX) is usually carried out by refluxing alcoholic solution of (IX) containing concentrated sulfuric acid. The

product is isolated by neutralisation of the reaction mixture and extraction. It can be appreciated that thus prepared (XI) is an ester.


The fifteenth aspect of the invention provides a compound of formula (XI)



(XI)

5


The sixteenth aspect of the invention provides a process for the manufacture of 1H-Pyrrolo[2,3-*b*]pyridine-5-carbonitrile (XI) comprising reaction of 5-bromo-1H-pyrrolo[2,3-*b*]pyridine with  $\text{Zn}(\text{CN})_2$  in the presence of a suitable palladium catalyst such as  $\text{Pd}(\text{PPh}_3)_4$ .



(XI)

10

The present invention encompasses one or more compounds as defined in the third, fifth, seventh, ninth, eleventh, thirteenth and fifteenth of the invention as set out below;



- The present invention also encompasses a process for manufacturing a compound of the first aspect, the process comprising providing a starting material, which is commercially available or can be produced by a method known in the art, converting the starting material to form an intermediate compound of the third, fifth, seventh, ninth, eleventh, thirteenth and fifteenth aspects using a process as described above or a process known in the art (and optionally converting the intermediate compound so formed into another intermediate compound) and then converting the intermediate compound into a compound of the first aspect using a process as described above or a process known in the art (and optionally converting the compound of the first aspect so formed into another compound of the first aspect).
- 15 The seventeenth aspect of the invention provides a composition comprising a compound according to the first aspect of the invention in combination with a pharmaceutically acceptable carrier, diluent or excipient.



The composition may also comprise one or more additional active agent, such as an anti-inflammatory agent (for example a p38 inhibitor, glutamate receptor antagonist, or a calcium channel antagonist), AMPA receptor antagonist, a chemotherapeutic agent and/or an antiproliferative agent.

5

Suitable carriers and/or diluents are well known in the art and include pharmaceutical grade starch, mannitol, lactose, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, (or other sugar), magnesium carbonate, gelatin, oil, alcohol, detergents, emulsifiers or water (preferably sterile). The composition may be a mixed preparation of a composition or may be a combined preparation for simultaneous, separate or sequential use (including administration).

The composition according to the invention for use in the aforementioned indications may be administered by any convenient method, for example by oral (including by inhalation), parenteral, mucosal (e.g. buccal, sublingual, nasal), rectal or transdermal administration and the compositions adapted accordingly.

20 For oral administration, the composition can be formulated as liquids or solids, for example solutions, syrups, suspensions or emulsions, tablets, capsules and lozenges.

25 A liquid formulation will generally consist of a suspension or solution of the compound or physiologically acceptable salt in a suitable aqueous or non-aqueous liquid carrier(s) for example water, ethanol, glycerine, polyethylene glycol or oil. The formulation may also contain a suspending agent, preservative, flavouring or colouring agent.

A composition in the form of a tablet can be prepared using any suitable pharmaceutical carrier(s) routinely used for preparing solid formulations. Examples of such carriers include magnesium stearate, starch, lactose, sucrose and microcrystalline cellulose.

5

A composition in the form of a capsule can be prepared using routine encapsulation procedures. For example, powders, granules or pellets containing the active ingredient can be prepared using standard carriers and then filled into a hard gelatine capsule; alternatively, a dispersion or suspension  
10 can be prepared using any suitable pharmaceutical carrier(s), for example aqueous gums, celluloses, silicates or oils and the dispersion or suspension then filled into a soft gelatine capsule.

Compositions for oral administration may be designed to protect the active  
15 ingredient against degradation as it passes through the alimentary tract, for example by an outer coating of the formulation on a tablet or capsule.

Typical parenteral compositions consist of a solution or suspension of the compound or physiologically acceptable salt in a sterile aqueous or non-  
20 aqueous carrier or parenterally acceptable oil, for example polyethylene glycol, polyvinyl pyrrolidone, lecithin, arachis oil or sesame oil. Alternatively, the solution can be lyophilised and then reconstituted with a suitable solvent just prior to administration.

25 Compositions for nasal or oral administration may conveniently be formulated as aerosols, drops, gels and powders. Aerosol formulations typically comprise a solution or fine suspension of the active substance in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the

form of a cartridge or refill for use with an atomising device. Alternatively the sealed container may be a unitary dispensing device such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve, which is intended for disposal once the contents of the container have been exhausted. Where the dosage form comprises an aerosol dispenser, it will contain a pharmaceutically acceptable propellant. The aerosol dosage forms can also take the form of a pump-atomiser.

Compositions suitable for buccal or sublingual administration include tablets, lozenges and pastilles, wherein the active ingredient is formulated with a carrier such as sugar and acacia, tragacanth, or gelatin and glycerin.

Compositions for rectal or vaginal administration are conveniently in the form of suppositories (containing a conventional suppository base such as cocoa butter), pessaries, vaginal tabs, foams or enemas.

Compositions suitable for transdermal administration include ointments, gels, patches and injections including powder injections.

Conveniently the composition is in unit dose form such as a tablet, capsule or ampoule.

The eighteenth aspect of the invention provides a process for the manufacture of a composition according to the seventeenth aspect of the invention. The manufacture can be carried out by standard techniques well known in the art and comprises combining a compound according to the first aspect of the invention and the pharmaceutically acceptable carrier or diluent and optionally one or more additional active agents. The composition may be in any form including a tablet, a liquid, a capsule, and a powder or in the form of a food

product, e.g. a functional food. In the latter case the food product itself may act as the pharmaceutically acceptable carrier.

5 The nineteenth aspect of the present invention relates to a compound of the first aspect, or a composition of the seventeenth aspect, for use in medicine.

The compounds of the present invention are inhibitors of JNK, such as JNK1, JNK2, or JNK3. In particular, the compounds of the present invention are inhibitors of JNK3. Preferably, the compounds of the present invention inhibit  
10 JNK3 selectively (i.e. the compounds of the invention preferably show greater activity against JNK3 than JNK1 and 2). For the purpose of this invention, an inhibitor is any compound, which reduces or prevents the activity of the JNK enzyme.

15 The compounds are therefore useful for conditions for which inhibition of JNK activity is beneficial. Thus, preferably, this aspect provides a compound of the first aspect, or a composition of the seventeenth aspect of the present invention, for the prevention or treatment of a JNK-mediated disorder. The compounds of the first aspect of the invention may thus be used for the  
20 inhibition of JNK, more preferably for the inhibition of JNK3.

A "JNK-mediated disorder" is any disease or deleterious condition in which JNK plays a role. Examples include neurodegenerative disorder (including dementia), inflammatory disease, a disorder linked to apoptosis, particularly  
25 neuronal apoptosis, autoimmune disease, destructive bone disorder, proliferative disorder, cancer, infectious disease, allergy, ischemia reperfusion injury, heart attack, angiogenic disorder, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin induced platelet aggregation and any condition

associated with prostaglandin endoperoxidase synthase-2. The compounds of the present invention may be used for any of these JNK-mediated disorders.

5 The compounds of the present invention are particularly useful for the prevention or treatment of a neurodegenerative disorder. In particular, the neurodegenerative disorder results from apoptosis and/or inflammation. Examples of neurodegenerative disorders are: dementia; Alzheimer's disease; Parkinson's disease; Amyotrophic Lateral Sclerosis; Huntington's disease; senile chorea; Sydenham's chorea; hypoglycemia; head and spinal cord trauma  
10 including traumatic head injury; acute and chronic pain; epilepsy and seizures; olivopontocerebellar dementia; neuronal cell death; hypoxia-related neurodegeneration; acute hypoxia; glutamate toxicity including glutamate neurotoxicity; cerebral ischemia; dementia linked to meningitis and/or neurosis; cerebrovascular dementia; or dementia in an HIV-infected patient.

15

The neurodegenerative disorder may be a peripheral neuropathy, including mononeuropathy, multiple mononeuropathy or polyneuropathy. Examples of peripheral neuropathy may be found in diabetes mellitus, Lyme disease or uremia; peripheral neuropathy caused by a toxic agent; demyelinating disease  
20 such as acute or chronic inflammatory polyneuropathy, leukodystrophies, or Guillain-Barré syndrome; multiple mononeuropathy secondary to a collagen vascular disorder (e.g. polyarteritis nodosa, SLE, Sjögren's syndrome); multiple mononeuropathy secondary to sarcoidosis; multiple mononeuropathy secondary to a metabolic disease (e.g. diabetes or amyloidosis); or multiple  
25 mononeuropathy secondary to an infectious disease (e.g. Lyme disease or HIV infection).

The compounds of the invention can also be used to prevent or treat disorders resulting from inflammation. These include, for example, inflammatory bowel



- disorder, bronchitis, asthma, acute pancreatitis, chronic pancreatitis, allergies of various types, and possibly Alzheimer's disease. Autoimmune diseases which may also be treated or prevented by the compounds of the present invention include rheumatoid arthritis, systemic lupus erythematosus, glomerulonephritis, scleroderma, chronic thyroiditis, Graves's disease, autoimmune gastritis, diabetes, autoimmune haemolytic anaemia, autoimmune neutropenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, ulcerative colitis, Crohn's disease, psoriasis or graft vs host disease.
- 10 A compound of the present invention may be administered simultaneously, subsequently or sequentially with one or more other active agent, such as an anti-inflammatory agent e.g. p38 inhibitor, AMPA receptor antagonist, glutamate receptor antagonist, calcium channel antagonist, a chemotherapeutic agent or an antiproliferative agent. For example, for acute treatment, a p38
- 15 inhibitor may be administered to a patient prior to administering a compound of the present invention.

The compounds of the invention will normally be administered in a daily dosage regimen (for an adult patient) of, for example, an oral dose of between 1

20 mg and 2000 mg, preferably between 30 mg and 1000 mg, e.g. between 10 and 250 mg or an intravenous, subcutaneous, or intramuscular dose of between 0.1 mg and 100 mg, preferably between 0.1 mg and 50 mg, e.g. between 1 and 25 mg of the compound of the formula (I) or a physiologically acceptable salt thereof calculated as the free base, the compound being administered 1 to 4

25 times per day. Suitably the compounds will be administered for a period of continuous therapy, for example for a week or more.

The twentieth aspect of the invention relates to a method of treating or preventing a JNK-mediated disorder in an individual, which method comprises

administering to said individual a compound of the first aspect or a composition of the seventeenth aspect. The active compound is preferably administered in a cumulative effective amount. The individual may be in need of the treatment or prevention. Any of the JNK-mediated disorders listed above in relation to the  
5 nineteenth aspect may be the subject of treatment or prevention according to the twentieth aspect. One or more other active agent may be administered to the individual simultaneously, subsequently or sequentially to administering the compound. The other active agent may be an anti-inflammatory agent such as a p38 inhibitor, glutamate receptor antagonist, AMPA receptor antagonist,  
10 calcium channel antagonist, a chemotherapeutic agent or an antiproliferative agent, but is preferably p38 inhibitor for acute treatment.

The twenty first aspect of the present invention provides the use of a compound of the first aspect in the manufacture of a medicament for the prevention or  
15 treatment of a JNK-mediated disorder. The medicament may be used for treatment or prevention of any of the JNK-mediated disorders listed above in relation to the nineteenth aspect. Again, the compound of the present invention may be administered simultaneously, subsequently or sequentially with one or more other active agent, preferably a p38 inhibitor for acute treatment.

20

In the twenty second aspect of the invention, there is provided an assay for determining the activity of the compounds of the present invention, comprising providing a system for assaying the activity and assaying the activity of the compound. Preferably the assay is for the JNK inhibiting activity of the  
25 compound, more preferably it is for the JNK3-specific inhibiting activity of the compounds. The compounds of the invention may be assayed in vitro, in vivo, in silico, or in a primary cell culture or a cell line. In vitro assays include assays that determine inhibition of either the kinase activity or ATPase activity of activated JNK. Alternatively, in vitro assays may quantitate the ability of a

compound to bind JNK and may be measured either by radiolabelling the compound prior to binding, then isolating the inhibitor/JNK complex and determining the amount of the radiolabel bound or by running a competition experiment where new inhibitors are incubated with JNK bound to known  
5 radioligands. An example of an assay, which may be used, is Scintillation Proximity Assay (SPA), preferably using radiolabelled ATP. Another example is ELISA. Any type or isoform of JNK may be used in these assays.


In the twenty third aspect, there is provided a method of inhibiting the activity  
10 or function of a JNK, particularly JNK3, which method comprises exposing a JNK to a compound or a composition of the first or seventeenth aspect of the present invention. The method may be performed in a research model, in vitro, in silico, or in vivo such as in an animal model. A suitable animal model may be a kainic acid model in rat or mice, traumatic brain injury model in rat, or  
15 MPTP in mice.

All features of each of the aspects apply to all other aspects *mutatis mutandis*.


The invention will now be illustrated by the following non-limiting examples.

## EXAMPLES

## 5 Synthesis of example inhibitor 8



## 1H-Pyrrolo[2,3-b]pyridine-5-carbonitrile (2)



- 10 A mixture of bromide 1 (10.0 g, 50.8 mmol), ZnCl<sub>2</sub> (3.58 g, 30.5 mmol), and Pd(PPh<sub>3</sub>)<sub>4</sub> (3.52 g, 3.05 mmol) in DMF (110 mL) was heated at 80 °C overnight. The solvent was evaporated and the residue separated by silicagel

5

### 1H-Pyrrolo[2,3-b]pyridine-5-carboxylic acid ethyl ester (3)



15

### 3-Iodo-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid ethyl ester (4)






To a solution of 3 (100 mg, 0.526 mmol) in DMF (1.3 mL) was added I<sub>2</sub> (162.56 mg, 0.64 mmol) followed by KOH (43.0 mg, 0.77 mmol). The reaction mixture was stirred for 35 min, and treated with a mixture of 0.1 M phosphate buffer (2.0 mL): saturated aqueous Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> (0.5 mL). The suspension was stirred at r.t. for 15 min. The solid precipitate was filtered off, washed with water (2.0 mL), and dried under high vacuum to give 4 as white solid (123.0 mg, 74%); <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ 1.34 (t, *J* = 7.1 Hz, 3H), 4.35 (q, *J* = 7.1 Hz, 2H), 7.87 (s, 1H), 8.15 (d, *J* = 2.0 Hz, 1H), 8.80 (d, *J* = 2.0 Hz, 1H), 12.50 (bs, 1H).

10


**1-Benzenesulfonyl-3-iodo-1H-pyrrolo[2,3-*b*]pyridine-5-carboxylic acid ethyl ester (5)**



To a suspension of 4 (112.5 mg, 0.356 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2.2 mL) was added benzenesulfonyl chloride (69.6 μL, 0.55 mmol), tetra-*n*-butylammonium hydrogen sulfate (14.9 mg, 0.044 mmol) and 50% aqueous NaOH (29 μL), and the reaction mixture was stirred for 1 h. The organic layer was separated and the residue was extracted with CH<sub>2</sub>Cl<sub>2</sub> (2x3 mL). The combined organic solutions were dried (MgSO<sub>4</sub>) and concentrated to give oil, which was triturated with methanol (1.0 mL) to give white solid. The solid was filtered off, washed with methanol (2x1 mL) and dried overnight in vacuum to give product 5 as a white solid (139.2 mg, 64%); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.42 (t, *J* = 7.1 Hz, 3H), 4.43 (q, *J* = 7.1 Hz, 2H), 7.49-7.53 (m, 2H), 7.59-7.64 (m, 1H), 7.94 (s, 1H), 8.21-8.25 (m, 2H), 8.31 (d, *J* = 2.0 Hz, 1H), 9.08 (d, *J* = 2.0 Hz, 1H).


25

**1-Benzenesulfonyl-3-furan-3-yl-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid ethyl ester (6)**



A mixture of 5 (100 mg, 0.219 mmol), EtOH (1.3 mL), toluene (1.3 mL), furan-3-boronic acid (37.4 mg, 0.33 mmol), 1M aq. Na<sub>2</sub>CO<sub>3</sub> (0.55 mL, 0.55 mmol), LiCl (28 mg, 0.66 mmol) and PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (12.9 mg, 18.4 μmol) was refluxed for 17 min. The organic layer was separated, brine was added, and the aqueous layer was extracted with AcOEt. The combined organic solutions were concentrated and separated by means of silicagel chromatography using hexane:CH<sub>2</sub>Cl<sub>2</sub> as eluent (in gradient up to 15% AcOEt) to give 6 as a tan solid (66.8 mg, 77%); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.42 (t, J = 7.1 Hz, 3H), 4.43 (q, J = 7.1 Hz, 2H), 6.71 (dd, J = 1.8, 0.9 Hz, 1H), 7.48-7.54 (m, 2H), 7.56 (t, J = 1.7 Hz, 1H), 7.58-7.63 (m, 1H), 7.85 (t, J = 0.9 Hz, 1H), 7.89 (s, 1H), 8.22-8.27 (m, 2H), 8.60 (d, J = 2.0 Hz, 1H), 9.12 (d, J = 2.0 Hz, 1H).

**3-Furan-3-yl-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid (7)**




To a suspension of 6 (65.5 mg, 0.165 mmol) in EtOH (1.0 mL), was added 10% aqueous NaOH (0.5 mL, about 1.25 mmol), and the reaction mixture was refluxed for 0.5 h. The mixture was cooled to r.t. and EtOH was evaporated in vacuum. The residual solution was treated with glacial acetic acid (75 μL, 1.25 mmol). The suspension which formed was stirred at r.t. for 30 min. The solid

was filtered off, washed with water, and dried under high vacuum to afford 7 as a tan powder (34.1 mg, 91%);  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  6.94 (d,  $J = 1.7$  Hz, 1H), 7.74 (t,  $J = 1.5$  Hz, 1H), 7.88 (d,  $J = 2.2$  Hz, 1H), 8.17 (s, 1H), 8.62 (d,  $J = 1.8$  Hz, 1H), 8.81 (d,  $J = 1.8$  Hz, 1H), 12.16 (bs, 1H).

5

**3-Furan-3-yl-1H-pyrrolo[2,3-b]pyridine-5-carboxylic acid (2-methoxy-phenyl)-amide (8)**



A mixture of 2-methoxy-phenylamine (21.55 mg, 0.175 mmol), carboxylic acid 7 (20 mg, 88  $\mu\text{mol}$ ), BOP (50.42 mg, 0.114 mmol), HOBT (17.8 mg, 0.132 mmol) and  $i\text{-Pr}_2\text{NEt}$  (30.7  $\mu\text{L}$ , 0.176 mmol) in dry DMF (0.6 mL) was stirred at r.t. for 2 h. Separation of the crude reaction mixture by LCMS (column LUNA 10  $\mu$  C18(2) 00G-4253-V0 250x50 mm) using water – acetonitrile (0.1% AcOH) as eluent (in gradient; flow 80 mL/min) afforded amide 8 (13.11 mg, 45%) as a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3:5\%\text{CD}_3\text{OD}$ )  $\delta$  3.93 (s, 3H), 6.71 (dd,  $J = 1.8, 0.8$  Hz, 1H), 6.94 (dd,  $J = 8.0, 1.5$  Hz, 1H), 7.03 (ddd,  $J = 7.9, 7.7, 1.5$  Hz, 1H), 7.10 (ddd,  $J = 8.0, 7.7, 1.7$  Hz, 1H), 7.50 (s, 1H), 7.53 (t,  $J = 1.7$  Hz, 1H), 7.84 (t,  $J = 1.1$  Hz, 1H), 8.48 (dd,  $J = 7.9, 1.7$  Hz, 1H), 8.61 (d,  $J = 2.0$  Hz, 1H), 8.81 (d,  $J = 2.0$  Hz, 1H); LCMS  $m/e$  334 ( $\text{M}+\text{H}$ ), 375 ( $\text{M}+\text{MeCN}+\text{H}$ ).

# Biological activity

## JNK1, JNK2, JNK3 - SPA assay

5

10

15

20

25

1. Compound is dissolved in DMSO to a convenient concentration and this is diluted in 10% DMSO to a five times concentrate of the desired starting concentration (frequently 1:100).
2. 10 µl of 500 mM EDTA is added to alternative wells of the Opti-plate row, which will receive kinase reaction plus DMSO. This creates the negative control.
3. For the JNK2 and JNK3 assay, compounds are prepared in six 2-fold dilutions with water and each concentration is tested in duplicate. For the JNK1 assay compounds are prepared in four 5-fold dilutions with water which are tested in triplicate. Controls are treated identically.
4. 20 µl per well of each compound concentration is transferred to an Opti-plate, in duplicate.
5. 30 µl (JNK2/3 SPA) or 50 µl (JNK1 SPA) of substrate solution (25 mM HEPES pH 7.5, 10mM magnesium acetate with 3.33µM ATP (JNK2/3) or 2µM ATP (JNK1), approximately 7.5 kBq [ $\gamma$ -<sup>33</sup>P] ATP, GST-c-Jun, in water) is added to each well.
6. 50 µl (JNK2/3 SPA) or 30 µl (JNK1 SPA) of kinase solution (JNK in 25 mM HEPES pH 7.5, 10mM Mg Acetate) is added to each well.

Kinase	Kinase per well (µg)	GST-c-Jun per well (µg)
--------	----------------------	-------------------------

JNK1	0.25	1
JNK2	0.2	1.2
JNK3	0.16	1.2

7. The plate is incubated for 30 minutes at room temperature.
8. 100  $\mu$ l of bead/stop solution is added to each well (5 mg/ml glutathione-PVT-SPA beads, 40 mM ATP in PBS).
9. Plates are sealed and incubated for 30 minutes at room temperature,  
5 centrifuged for 10 minutes at 2500g and counted.
10. The  $IC_{50}$  values are calculated as the concentration of the compound being tested at which the phosphorylation of c-Jun is decreased to 50% of the control value. Example  $IC_{50}$  values for the compounds of this invention are given in Table 1.





10

### p38 ELISA

Active p38 kinase (100 ng; Upstate) was added to 2  $\mu$ g GST-ATF2 substrate (NEB) in 250 mM Hepes pH 7.5/100 mM MgAc/50  $\mu$ M ATP (final) in the  
15 presence or absence of compounds in 50 $\mu$ l. The mixture was incubated at 30°C for 1 hour, and then diluted with 200  $\mu$ l PBS-Tween (0.05 %). From this, duplicate volumes of 100  $\mu$ l were added to a Reacti-Bind glutathione coated plate (Pierce) and incubated for 1 hour. After washing 3 times with PBS-Tween (0.05 %), rabbit anti-phospho-ATF2 (Thr71) antibody (NEB) was added  
20 at 1:500, and incubated for another hour at room temperature. After 3 additional washes with PBS-Tween (0.05 %), 100  $\mu$ l of anti-rabbit IgG alkaline phosphatase-conjugated secondary antibody (Sigma) was added at 1:1000, the reaction was incubated for a further hour, washed 3 times, and then phosphatase

substrate (Sigma) was added (100  $\mu$ l per well; 3 tablets in 5 ml water). After incubation in the dark at 37°C for 1 hour, the reaction mixture was transferred to a clear 96 well plate, and the absorbance at 405 nm was read. The IC<sub>50</sub> values are calculated as the concentration of the compound being tested at which the phosphorylation of ATF2 is decreased to 50% of the control value. Example IC<sub>50</sub> values for the compounds of this invention are given in Table 1 (last column).


Table 1. IC<sub>50</sub> values for selected compounds against JNK1, JNK2, JNK3, and p38 MAP kinase

Compound	JNK3 IC <sub>50</sub> (nM)
	<500
	<500
	<1000
	<500




## CLAIMS

1. A compound of formula (I):



- 5 and the pharmaceutically acceptable salts, and other pharmaceutically acceptable biohydrolyzable derivatives thereof;

wherein  $R^1$  is an optionally substituted  $C_{3-12}$  carbocyclyl or  $C_{3-12}$  heterocyclyl group or group of formula (II)



10

wherein X is  $NR^3$ , O, S or  $(CR^{22}R^{22})_n$ , Y is absent or is  $NR^{23}$ , O, or  $(CR^{23}R^{23})_n$ ,  $R^2$  is optionally substituted  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{3-12}$  carbocyclyl or  $C_{3-12}$  heterocyclyl, and  $R^4$  is an optionally substituted five or six membered heterocyclyl group or an optionally substituted six membered

15 carbocyclyl group.

2. A compound as claimed in claim 1 wherein the optionally substituted carbocyclyl or heterocyclyl group of  $R^1$  is optionally fused to a partially saturated, unsaturated or fully saturated five to seven membered ring containing
- 20 zero to three heteroatoms, and each substitutable carbon atom in  $R^1$ , including the optional fused ring, is optionally and independently substituted by one or more of halogen,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl, halo $C_{1-12}$ alkyl,  $C_{3-12}$  carbocyclyl,  $C_{3-12}$  heterocyclyl,  $(CH_2)_nOR^5$ ,  $(CH_2)_nNR^5_2(CH_2)_nSR^5$ ,  $OR^5$ ,  $SR^5$ ,  $NO_2$ ,  $CN$ ,  $NR^5_2$ ,  $NR^5COR^5$ ,  $NR^5CONR^5_2$ ,  $NR^5COR^5$ ,  $NR^5CO_2R^5$ ,  $CO_2R^5$ ,
- 25  $COR^5$ ,  $CONR^5_2$ ,  $S(O)_2R^5$ ,  $SONR^5_2$ ,  $S(O)R^5$ ,  $SO_2NR^5_2$ , or  $NR^5S(O)_2R^5$  wherein

the C<sub>1-12</sub> alkyl group optionally contains one or more insertions selected from -O-, -N(R<sup>5</sup>)- -S-, -S(O)- and -S(O<sub>2</sub>)-; and each saturated carbon in the optional fused ring is further optionally and independently substituted by =O, =S, NNR<sup>6</sup><sub>2</sub>, =N-OR<sup>6</sup>, =NNR<sup>6</sup>COR<sup>6</sup>, =NNR<sup>6</sup>CO<sub>2</sub>R<sup>6</sup>, =NNSO<sub>2</sub>R<sup>6</sup>, or =NR<sup>6</sup>; and each  
 5 substitutable nitrogen atom in R<sup>1</sup> is optionally substituted by R<sup>7</sup>, COR<sup>7</sup>, SO<sub>2</sub>R<sup>7</sup> or CO<sub>2</sub>R<sup>7</sup>;

wherein n is 1 to 6, preferably n is 1, 2 or 3;

wherein R<sup>5</sup> is hydrogen, C<sub>1-12</sub> alkyl, C<sub>3-12</sub> carbocyclyl or C<sub>3-12</sub> heterocyclyl,  
 10 optionally substituted by one or more of C<sub>1-6</sub> alkyl, C<sub>3-12</sub> carbocyclyl, C<sub>3-12</sub> heterocyclyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>8</sup>, SR<sup>8</sup>, NO<sub>2</sub>, CN, NR<sup>8</sup>R<sup>8</sup>, NR<sup>8</sup>COR<sup>8</sup>, NR<sup>8</sup>CONR<sup>8</sup>R<sup>8</sup>, NR<sup>8</sup>COR<sup>8</sup>, NR<sup>8</sup>CO<sub>2</sub>R<sup>8</sup>, CO<sub>2</sub>R<sup>8</sup>, COR<sup>8</sup>, CONR<sup>8</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>8</sup>, SONR<sup>8</sup><sub>2</sub>, S(O)R<sup>8</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>8</sup>, NR<sup>8</sup>S(O)<sub>2</sub>R<sup>8</sup>, wherein the C<sub>1-12</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting  
 15 of -O-, -N(R<sup>8</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>8</sup> may be the same or different and is as defined below;

wherein two R<sup>5</sup> in NR<sup>5</sup><sub>2</sub> may optionally form a partially saturated, unsaturated or fully saturated three to seven membered ring containing one to three  
 20 heteroatoms, optionally and independently substituted by one or more of C<sub>1-6</sub> alkyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>8</sup>, SR<sup>8</sup>, NO<sub>2</sub>, CN, NR<sup>8</sup>R<sup>8</sup>, NR<sup>8</sup>COR<sup>8</sup>, NR<sup>8</sup>CONR<sup>8</sup>R<sup>8</sup>, NR<sup>8</sup>COR<sup>8</sup>, NR<sup>8</sup>CO<sub>2</sub>R<sup>8</sup>, CO<sub>2</sub>R<sup>8</sup>, COR<sup>8</sup>, CONR<sup>8</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>8</sup>, SONR<sup>8</sup><sub>2</sub>, S(O)R<sup>8</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>8</sup>, NR<sup>8</sup>S(O)<sub>2</sub>R<sup>8</sup>,

wherein the C<sub>1-6</sub> alkyl group optionally incorporates one or two insertions  
 25 selected from the group consisting of -O-, -N(R<sup>8</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>8</sup> may be the same or different and is as defined below;

wherein R<sup>6</sup> is hydrogen, C<sub>1-12</sub> alkyl, C<sub>3-12</sub> carbocyclyl or C<sub>3-12</sub> heterocyclyl, optionally substituted by one or more of C<sub>1-6</sub> alkyl, halogen, C<sub>1-6</sub> haloalkyl,

OR<sup>8</sup>, SR<sup>8</sup>, NO<sub>2</sub>, CN, NR<sup>8</sup>R<sup>8</sup>, NR<sup>8</sup>COR<sup>8</sup>, NR<sup>8</sup>CONR<sup>8</sup>R<sup>8</sup>, NR<sup>8</sup>COR<sup>8</sup>, NR<sup>8</sup>CO<sub>2</sub>R<sup>8</sup>, CO<sub>2</sub>R<sup>8</sup>, COR<sup>8</sup>, CONR<sup>8</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>8</sup>, S(O)R<sup>8</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>8</sup>, NR<sup>8</sup>S(O)<sub>2</sub>R<sup>8</sup>, wherein the C<sub>1-12</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -N(R<sup>8</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>8</sup> may  
 5 be the same or different and is as defined below;

wherein R<sup>7</sup> is hydrogen, C<sub>6-12</sub> aryl, C<sub>1-6</sub> alkyl or C<sub>1-6</sub> haloalkyl;

wherein R<sup>8</sup> is hydrogen, C<sub>1-6</sub> alkyl, or C<sub>1-6</sub> haloalkyl.

10

3. A compound as claimed in claim 1 or 2 wherein Y is absent or is NR<sup>23</sup>, O, (CR<sup>23</sup>R<sup>23</sup>)<sub>n</sub>,

wherein each R<sup>23</sup> is H, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy or C<sub>1-4</sub> haloalkyl;

and n is 1 to 6, preferably n is 1, 2, 3 or 4; and

15

R<sup>2</sup> is optionally substituted C<sub>1-12</sub> alkyl, C<sub>2-12</sub> alkenyl, C<sub>2-12</sub> alkynyl, C<sub>3-12</sub> carbocyclyl or C<sub>3-12</sub> heterocyclyl, wherein the optionally substituted carbocyclyl or heterocyclyl group is optionally fused to one to three unsaturated, partially unsaturated or fully saturated five to seven membered

20 rings containing zero to three heteroatoms;

each substitutable carbon atom in R<sup>2</sup>, including the optional fused ring, is optionally and independently substituted by one or more of C<sub>1-12</sub> alkyl, C<sub>3-12</sub> cycloalkyl, C<sub>3-12</sub> heterocycloalkyl, C<sub>3-12</sub> aryl, C<sub>3-12</sub> heteroaryl halogen, C<sub>1-12</sub> haloalkyl, OR<sup>9</sup>, SR<sup>9</sup>, NO<sub>2</sub>, CN, NR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>COR<sup>9</sup>, NR<sup>9</sup>CONR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>COR<sup>9</sup>, NR<sup>9</sup>CO<sub>2</sub>R<sup>9</sup>, CO<sub>2</sub>R<sup>9</sup>, COR<sup>9</sup>, CONR<sup>9</sup>R<sup>9</sup>, S(O)<sub>2</sub>R<sup>9</sup>, SONH<sub>2</sub>, S(O)R<sup>9</sup>, SO<sub>2</sub>NR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>S(O)<sub>2</sub>R<sup>9</sup>, wherein each R<sup>9</sup> may be the same or different and is as defined  
 25 below and wherein:

- the C<sub>1-12</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -C(O)-, -N(R<sup>9</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>9</sup> may be the same or different and is as defined above;
- the C<sub>1-12</sub> alkyl, C<sub>3-12</sub> cycloalkyl, C<sub>3-12</sub> heterocycloalkyl, C<sub>3-12</sub> aryl, or C<sub>3-12</sub> heteroaryl groups are optionally substituted by one or more of halogen, C<sub>1-12</sub> haloalkyl, OR<sup>9</sup>, SR<sup>9</sup>, NO<sub>2</sub>, CN, NR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>COR<sup>9</sup>, NR<sup>9</sup>CONR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>COR<sup>9</sup>, NR<sup>9</sup>CO<sub>2</sub>R<sup>9</sup>, CO<sub>2</sub>R<sup>9</sup>, COR<sup>9</sup>, CONR<sup>9</sup>R<sup>9</sup>, S(O)<sub>2</sub>R<sup>9</sup>, SONH<sub>2</sub>, S(O)R<sup>9</sup>, SO<sub>2</sub>NR<sup>9</sup>R<sup>9</sup>, NR<sup>9</sup>S(O)<sub>2</sub>R<sup>9</sup>, wherein each R<sup>9</sup> may be the same or different and is as defined below; and
- 10 the C<sub>3-12</sub> cycloalkyl, C<sub>3-12</sub> heterocycloalkyl, C<sub>3-12</sub> aryl, or C<sub>3-12</sub> heteroaryl groups are optionally substituted by one or more C<sub>1-12</sub> alkyl groups;
- each saturated carbon in R<sup>2</sup>, including the optional fused ring, is further optionally and independently substituted by =O, =S, NNR<sup>9</sup>R<sup>9</sup>, =N-OR<sup>9</sup>, =NNHCOR<sup>9</sup>, =NNHCO<sub>2</sub>R<sup>9</sup>, =NNSO<sub>2</sub>R<sup>9</sup>, or =NR<sup>9</sup>, wherein each R<sup>9</sup> may be the
- 15 same or different and is as defined below; and
- each substitutable nitrogen atom in R<sup>2</sup> is optionally substituted by R<sup>10</sup>, COR<sup>9</sup>, SO<sub>2</sub>R<sup>9</sup> or CO<sub>2</sub>R<sup>9</sup> wherein each R<sup>9</sup> and R<sup>10</sup> may be the same or different and is as defined below;
- wherein two R<sup>9</sup> in NR<sup>9</sup><sub>2</sub> may optionally form a partially saturated, unsaturated
- 20 or fully saturated three to seven membered ring containing one to three heteroatoms, optionally and independently substituted by one or more of C<sub>1-6</sub> alkyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>11</sup>, SR<sup>11</sup>, NO<sub>2</sub>, CN, NR<sup>11</sup>R<sup>11</sup>, NR<sup>11</sup>COR<sup>11</sup>, NR<sup>11</sup>CONR<sup>11</sup>R<sup>11</sup>, NR<sup>11</sup>COR<sup>11</sup>, NR<sup>11</sup>CO<sub>2</sub>R<sup>11</sup>, CO<sub>2</sub>R<sup>11</sup>, COR<sup>11</sup>, CONR<sup>11</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>11</sup>, SONR<sup>11</sup><sub>2</sub>, S(O)R<sup>11</sup>, SO<sub>2</sub>NR<sup>11</sup>R<sup>11</sup>, NR<sup>11</sup>S(O)<sub>2</sub>R<sup>11</sup>,
- 25 wherein the C<sub>1-6</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -N(R<sup>11</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>11</sup> may be the same or different and is as defined below;
- wherein R<sup>11</sup> is hydrogen, C<sub>1-6</sub> alkyl, or C<sub>1-6</sub> haloalkyl;

wherein  $R^9$  is hydrogen,  $C_{1-12}$  alkyl or  $C_{3-12}$  aryl, optionally substituted by one or more of  $C_{1-4}$  alkyl, halogen,  $C_{1-4}$  haloalkyl,  $OR^{12}$ ,  $SR^{12}$ ,  $NO_2$ ,  $CN$ ,  $NR^{12}R^{12}$ ,  $NR^{12}COR^{12}$ ,  $NR^{12}CONR^{12}R^{12}$ ,  $NR^{12}COR^{12}$ ,  $NR^{12}CO_2R^{12}$ ,  $CO_2R^{12}$ ,  $COR^{12}$ ,  $CONR^{12}_2$ ,  $S(O)_2R^{12}$ ,  $SONH_2$ ,  $S(O)R^{12}$ ,  $SO_2NR^{12}R^{12}$ ,  $NR^{12}S(O)_2R^{12}$ , wherein  
 5 the  $C_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-O-$ ,  $-N(R^{12})-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^{12}$  may be the same or different and is as defined below;

wherein  $R^{10}$  is  $C_{1-12}$  alkyl or  $C_{3-12}$  aryl, optionally substituted by one or more of  
 10  $C_{1-4}$  alkyl, halogen,  $C_{1-4}$  haloalkyl,  $OR^{12}$ ,  $SR^{12}$ ,  $NO_2$ ,  $CN$ ,  $NR^{12}R^{12}$ ,  $NR^{12}COR^{12}$ ,  $NR^{12}CONR^{12}R^{12}$ ,  $NR^{12}COR^{12}$ ,  $NR^{12}CO_2R^{12}$ ,  $CO_2R^{12}$ ,  $COR^{12}$ ,  $CONR^{12}_2$ ,  $S(O)_2R^{12}$ ,  $SONH_2$ ,  $S(O)R^{12}$ ,  $SO_2NR^{12}R^{12}$ ,  $NR^{12}S(O)_2R^{12}$ , wherein the  $C_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-O-$ ,  $-N(R^{12})-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^{12}$  may be the  
 15 same or different and is as defined below;

wherein  $R^{12}$  is hydrogen,  $C_{1-4}$  alkyl, or  $C_{1-4}$  haloalkyl.

4. A compound as claimed in any one of claims 1 to 3 wherein  $X$  is  $NR^3$ ;  
 20  $O$ ,  $S$  or  $(CR^{22}-R^{22})_n$  wherein  $R^{22}$  is independently one or more of halogen,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{1-12}$  haloalkyl,  $C_{6-12}$  carbocyclyl,  $C_{5-12}$  heterocyclyl,  $(CH_2)_nOR^5$ ,  $(CH_2)_nNR^5_2$ ,  $OR^5$ ,  $SR^5$ ,  $NO_2$ ,  $CN$ ,  $NR^5_2$ ,  $NR^5COR^5$ ,  $NR^5CONR^5_2$ ,  $NR^5COR^5$ ,  $NR^5CO_2R^5$ ,  $CO_2R^5$ ,  $COR^5$ ,  $CONR^5_2$ ,  $S(O)_2R^5$ ,  $SONR^5_2$ ,  $S(O)R^5$ ,  $SO_2NR^5_2$ , or  $NR^5S(O)_2R^5$  wherein each  $R^5$  may be the same  
 25 or different and is as defined above; and  
 wherein  $n$  is 1 to 6, preferably  $n$  is 1, 2, 3 or 4;



and wherein  $R^3$  is a lone electron pair, hydrogen,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{3-12}$  carbocyclyl or  $C_{3-12}$  heterocyclyl, each of which is optionally substituted, wherein:

the optionally substituted carbocyclyl or heterocyclyl group is optionally fused  
5 to one to three unsaturated, partially unsaturated or fully saturated five to seven membered rings containing zero to three heteroatoms,

each substitutable carbon atom in  $R^3$ , including the optional fused ring, is optionally and independently substituted by one or more of  $C_{1-12}$  alkyl,  $C_{3-12}$  cycloalkyl,  $C_{3-12}$  heterocycloalkyl,  $C_{3-12}$  aryl,  $C_{3-12}$  heteroaryl halogen,  $C_{1-12}$   
10 haloalkyl,  $OR^{13}$ ,  $SR^{13}$ ,  $NO_2$ ,  $CN$ ,  $NR^{13}R^{13}$ ,  $NR^{13}COR^{13}$ ,  $NR^{13}CONR^{13}R^{13}$ ,  $NR^{13}COR^{13}$ ,  $NR^{13}CO_2R^{13}$ ,  $CO_2R^{13}$ ,  $COR^{13}$ ,  $CONR^{13}R^{13}$ ,  $S(O)_2R^{13}$ ,  $SONH_2$ ,  $S(O)R^{13}$ ,  $SO_2NR^{13}R^{13}$ ,  $NR^{13}S(O)_2R^{13}$ , wherein each  $R^{13}$  may be the same or different and is as defined above and wherein:

the  $C_{1-12}$  alkyl group optionally incorporates one or two insertions selected from  
15 the group consisting of  $-O-$ ,  $-C(O)-$ ,  $-N(R^{13})-$ ,  $-S(O)-$  and  $-S(O)_2-$ , wherein each  $R^{13}$  may be the same or different and is as defined above;

the  $C_{1-12}$  alkyl,  $C_{3-12}$  cycloalkyl,  $C_{3-12}$  heterocycloalkyl,  $C_{3-12}$  aryl, or  $C_{3-12}$  heteroaryl groups are optionally substituted by one or more of halogen,  $C_{1-12}$  haloalkyl,  $OR^{13}$ ,  $SR^{13}$ ,  $NO_2$ ,  $CN$ ,  $NR^{13}R^{13}$ ,  $NR^{13}COR^{13}$ ,  $NR^{13}CONR^{13}R^{13}$ ,  
20  $NR^{13}COR^{13}$ ,  $NR^{13}CO_2R^{13}$ ,  $CO_2R^{13}$ ,  $COR^{13}$ ,  $CONR^{13}R^{13}$ ,  $S(O)_2R^{13}$ ,  $SONH_2$ ,  $S(O)R^{13}$ ,  $SO_2NR^{13}R^{13}$ ,  $NR^{13}S(O)_2R^{13}$ , wherein each  $R^{13}$  may be the same or different and is as defined below; and

the  $C_{3-12}$  cycloalkyl,  $C_{3-12}$  heterocycloalkyl,  $C_{3-12}$  aryl, or  $C_{3-12}$  heteroaryl groups are optionally substituted by one or more  $C_{1-12}$  alkyl groups;

25 each saturated carbon in  $R^2$ , including the optional fused ring, is further optionally and independently substituted by  $=O$ ,  $=S$ ,  $NNR^{13}R^{13}$ ,  $=N-OR^{13}$ ,  $=NNHCOR^{13}$ ,  $=NNHCO_2R^{13}$ ,  $=NNSO_2R^{13}$ , or  $=NR^{13}$ , wherein each  $R^{13}$  may be the same or different and is as defined below; and



each substitutable nitrogen atom in  $R^3$  is optionally substituted by  $R^{14}$ ,  $COR^{13}$ ,  $SO_2R^{13}$  or  $CO_2R^{13}$  wherein each  $R^{13}$  and  $R^{14}$  may be the same or different and is as defined below;

5 wherein two  $R^{13}$  in  $NR^{13}_2$  may optionally form a partially saturated, unsaturated or fully saturated three to seven membered ring containing one to three heteroatoms, optionally and independently substituted by one or more of  $C_{1-6}$  alkyl, halogen,  $C_{1-6}$  haloalkyl,  $OR^{15}$ ,  $SR^{15}$ ,  $NO_2$ ,  $CN$ ,  $NR^{15}R^{15}$ ,  $NR^{15}COR^{15}$ ,  $NR^{15}CONR^{15}R^{15}$ ,  $NR^{15}COR^{15}$ ,  $NR^{15}CO_2R^{15}$ ,  $CO_2R^{15}$ ,  $COR^{15}$ ,  $CONR^{15}_2$ ,  $S(O)_2R^{15}$ ,  $SONR^{15}_2$ ,  $S(O)R^{15}$ ,  $SO_2NR^{15}R^{15}$ ,  $NR^{15}S(O)_2R^{15}$ ,

10 wherein the  $C_{1-6}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-O-$ ,  $-N(R^{15})-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^{15}$  may be the same or different and is as defined below; wherein  $R^{15}$  is hydrogen,  $C_{1-6}$  alkyl, or  $C_{1-6}$  haloalkyl;

15 wherein  $R^{13}$  is hydrogen,  $C_{1-12}$  alkyl or  $C_{3-12}$  aryl, optionally substituted by one or more of  $C_{1-4}$  alkyl, halogen,  $C_{1-4}$  haloalkyl,  $OR^{16}$ ,  $SR^{16}$ ,  $NO_2$ ,  $CN$ ,  $NR^{16}R^{16}$ ,  $NR^{16}COR^{16}$ ,  $NR^{16}CONR^{16}R^{16}$ ,  $NR^{16}COR^{16}$ ,  $NR^{16}CO_2R^{16}$ ,  $CO_2R^{16}$ ,  $COR^{16}$ ,  $CONR^{16}_2$ ,  $S(O)_2R^{16}$ ,  $SONH_2$ ,  $S(O)R^{16}$ ,  $SO_2NR^{16}R^{16}$ ,  $NR^{16}S(O)_2R^{16}$ , wherein the  $C_{1-12}$  alkyl group optionally incorporates one or two insertions selected from  
20 the group consisting of  $-O-$ ,  $-N(R^{16})-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^{16}$  may be the same or different and is as defined below;

wherein  $R^{14}$  is  $C_{1-12}$  alkyl or  $C_{3-12}$  aryl, optionally substituted by one or more of  $C_{1-4}$  alkyl, halogen,  $C_{1-4}$  haloalkyl,  $OR^{16}$ ,  $SR^{16}$ ,  $NO_2$ ,  $CN$ ,  $NR^{16}R^{16}$ ,  $NR^{16}COR^{16}$ ,  
25  $NR^{16}CONR^{16}R^{16}$ ,  $NR^{16}COR^{16}$ ,  $NR^{16}CO_2R^{16}$ ,  $CO_2R^{16}$ ,  $COR^{16}$ ,  $CONR^{16}_2$ ,  $S(O)_2R^{16}$ ,  $SONH_2$ ,  $S(O)R^{16}$ ,  $SO_2NR^{16}R^{16}$ ,  $NR^{16}S(O)_2R^{16}$ , wherein the  $C_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-O-$ ,  $-N(R^{16})-$ ,  $-S(O)-$  and  $-S(O_2)-$ , wherein each  $R^{16}$  may be the same or different and is as defined below;

wherein  $R^{16}$  is hydrogen,  $C_{1-4}$  alkyl, or  $C_{1-4}$  haloalkyl;

wherein when X is  $NR^2$ ,  $R^2$  and  $R^3$  can form a 3 to 12 membered heterocycl  
 5 ring, more preferably a 5, 6, 7, 8, 9, 10, 11 or 12 membered ring, wherein said  
 ring can be partially saturated, unsaturated or fully saturated containing one to  
 three heteroatoms; wherein the heterocyclic group formed by  $R^2$  and  $R^3$  can  
 be optionally fused to one to three unsaturated, partially saturated or fully  
 saturated 5 to 7 membered rings containing zero to three heteroatoms, any of  
 10 said rings being optionally and independently substituted with one or more of  
 $C_{1-6}$  alkyl, halogen,  $C_{1-6}$  haloalkyl,  $OR^{22}$ ,  $SR^{22}$ ,  $NO_2$ ,  $CN$ ,  $NR^{22}R^{22}$ ,  $NR^{22}COR^{22}$ ,  
 $NR^{22}CONR^{22}R^{22}$ ,  $NR^{22}COR^{22}$ ,  $NR^{22}CO_2R^{22}$ ,  $CO_2R^{22}$ ,  $COR^{22}$ ,  $CONR^{22}_2$ ,  
 $S(O)_2R^{22}$ ,  $SONR^{22}_2$ ,  $S(O)R^{22}$ ,  $SO_2NR^{22}R^{22}$ ,  $NR^{22}S(O)_2R^{22}$ , wherein the  $C_{1-6}$   
 alkyl group optionally incorporates one or two insertions from  $-O-$ ,  $-N(R^{22})-$ ,  $-$   
 15  $S(O)-$  and  $-S(O)_2-$  and wherein each  $R^{22}$  may be the same or different.

5. A compound as claimed in any one of claims 1 to 4 wherein  $R^4$  is a six-  
 membered carbocyclyl group or a five or six-membered heterocyclyl group  
 containing from 1 to 4 heteroatoms independently selected from N, S or O,  
 20 wherein the optionally substituted six-membered carbocyclyl or five or six-  
 membered heterocyclyl group is optionally fused to a partially saturated,  
 unsaturated or fully saturated five to seven membered ring containing zero to  
 three heteroatoms, and each substitutable carbon or hetero-atom in  $R^4$  including  
 the optional fused ring, is optionally and independently substituted by one or  
 25 more of halogen,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{1-12}$  haloalkyl,  $C_{3-12}$   
 carbocyclyl,  $C_{3-12}$  heterocyclyl,  $(CH_2)_nOR^{17}$ ,  $(CH_2)_nNR^{17}_2$ ,  $OR^{17}$ ,  $SR^{17}$ ,  $NO_2$ ,  
 $CN$ ,  $NR^{17}_2$ ,  $NR^{17}COR^{17}$ ,  $NR^{17}CONR^{17}_2$ ,  $NR^{17}COR^{17}$ ,  $NR^{17}CO_2R^{17}$ ,  $CO_2R^{17}$ ,  
 $COR^{17}$ ,  $CONR^{17}_2$ ,  $S(O)_2R^{17}$ ,  $SONR^{17}_2$ ,  $S(O)R^{17}$ ,  $SO_2NR^{17}_2$ , or  $NR^{17}S(O)_2R^{17}$ ,  
 wherein the  $C_{1-12}$  alkyl group optionally contains one or more insertions

selected from -O-, -N(R<sup>12</sup>)-, -S-, -S(O)- and -S(O<sub>2</sub>)-; and each saturated carbon in the optional fused ring is further optionally and independently substituted by =O, =S, NNR<sup>18</sup><sub>2</sub>, =N-OR<sup>18</sup>, =NNR<sup>18</sup>COR<sup>18</sup>, =NNR<sup>18</sup>CO<sub>2</sub>R<sup>18</sup>, =NNSO<sub>2</sub>R<sup>18</sup>, or =NR<sup>18</sup>; and each substitutable nitrogen atom in R<sup>4</sup> is optionally substituted by  
 5 R<sup>19</sup>, COR<sup>19</sup>, SO<sub>2</sub>R<sup>19</sup> or CO<sub>2</sub>R<sup>19</sup>; wherein n is 1 to 6, preferably n is 1, 2 or 3; preferably, wherein each substitutable carbon or hetero-atom in R<sup>4</sup> is optionally and independently substituted by one or more of C<sub>1-6</sub> alkyl, OR<sup>20</sup>, SR<sup>20</sup>, NO<sub>2</sub>, CN, NR<sup>20</sup><sub>2</sub>, NR<sup>20</sup>COR<sup>20</sup>, NR<sup>20</sup>CONR<sup>20</sup><sub>2</sub>, NR<sup>20</sup>COR<sup>20</sup>, NHCO<sub>2</sub>R<sup>20</sup>, CO<sub>2</sub>R<sup>20</sup>, COR<sup>20</sup>, CONR<sup>20</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>20</sup>, SONR<sup>20</sup><sub>2</sub>, S(O)R<sup>20</sup>, SO<sub>2</sub>NR<sup>20</sup><sub>2</sub>, or NR<sup>20</sup>S(O)<sub>2</sub>R<sup>20</sup>;

10

wherein R<sup>20</sup> is hydrogen, C<sub>1-6</sub> alkyl, or C<sub>1-6</sub> haloalkyl;

wherein R<sup>17</sup> is hydrogen, C<sub>1-12</sub> alkyl, C<sub>3-12</sub> carbocyclyl or C<sub>3-12</sub> heterocyclyl, optionally substituted by one or more of C<sub>1-6</sub> alkyl, C<sub>3-12</sub> carbocyclyl, C<sub>3-12</sub>  
 15 heterocyclyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>21</sup>, SR<sup>21</sup>, NO<sub>2</sub>, CN, NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CONR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CO<sub>2</sub>R<sup>21</sup>, CO<sub>2</sub>R<sup>21</sup>, COR<sup>21</sup>, CONR<sup>21</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>21</sup>, SONR<sup>21</sup><sub>2</sub>, S(O)R<sup>21</sup>, SO<sub>2</sub>NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>S(O)<sub>2</sub>R<sup>21</sup>, wherein the C<sub>1-12</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -N(R<sup>21</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>21</sup>  
 20 may be the same or different and is as defined below;

wherein two R<sup>17</sup> in NR<sup>17</sup><sub>2</sub> may optionally form a partially saturated, unsaturated or fully saturated three to seven membered ring containing one to three heteroatoms, optionally and independently substituted by one or more of C<sub>1-6</sub>  
 25 alkyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>21</sup>, SR<sup>21</sup>, NO<sub>2</sub>, CN, NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CONR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CO<sub>2</sub>R<sup>21</sup>, CO<sub>2</sub>R<sup>21</sup>, COR<sup>21</sup>, CONR<sup>21</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>21</sup>, SONR<sup>21</sup><sub>2</sub>, S(O)R<sup>21</sup>, SO<sub>2</sub>NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>S(O)<sub>2</sub>R<sup>21</sup>, wherein the C<sub>1-6</sub> alkyl group optionally incorporates one or two insertions selected from the

group consisting of -O-, -N(R<sup>21</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>21</sup> may be the same or different and is as defined below;

wherein R<sup>18</sup> is hydrogen, C<sub>1-12</sub> alkyl, C<sub>3-12</sub> carbocyclyl or C<sub>3-12</sub> heterocyclyl,  
 5 optionally substituted by one or more of C<sub>1-6</sub> alkyl, halogen, C<sub>1-6</sub> haloalkyl, OR<sup>21</sup>, SR<sup>21</sup>, NO<sub>2</sub>, CN, NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CONR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>COR<sup>21</sup>, NR<sup>21</sup>CO<sub>2</sub>R<sup>21</sup>, CO<sub>2</sub>R<sup>21</sup>, COR<sup>21</sup>, CONR<sup>21</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>21</sup>, S(O)R<sup>21</sup>, SO<sub>2</sub>NR<sup>21</sup>R<sup>21</sup>, NR<sup>21</sup>S(O)<sub>2</sub>R<sup>21</sup>, wherein the C<sub>1-12</sub> alkyl group optionally incorporates one or two  
 10 S(O<sub>2</sub>)-, wherein each R<sup>21</sup> may be the same or different and is as defined below;

wherein R<sup>19</sup> is hydrogen, C<sub>6-12</sub> aryl, C<sub>1-6</sub> alkyl or C<sub>1-6</sub> haloalkyl;  
 wherein R<sup>21</sup> is hydrogen, C<sub>1-6</sub> alkyl, or C<sub>1-6</sub> haloalkyl.

15 6. A compound as claimed in any one of claims 1 to 5 wherein R<sup>1</sup> is an optionally substituted five or six membered carbocyclyl or heterocyclyl group selected from optionally substituted phenyl, acridine, benzimidazole, benzofuran, benzothiophene, benzoxazole, benzothiazole, cyclohexyl furan, imidazole, indole, isoindole, isoquinoline, isoxazole, isothiazole, morpholine,  
 20 naphthalene, oxazole, phenazine, phenothiazine, phenoxazine, piperazine, piperidine, pyrazole, pyridazine, pyridine, pyrrole, quinoline, quinolizine, tetrahydrofuran, tetrazine, tetrazole, thiophene, thiazole, thiomorpholine, thianaphthalene, thiopyran, triazine, triazole or trithiane.

25 7. A compound as claimed in any one of claims 1 to 6 wherein R<sup>1</sup> is a group of formula (II), wherein X is a group NR<sup>3</sup>, Y is absent and one or more of R<sup>2</sup> and R<sup>3</sup> is hydrogen, alkyl or cycloalkyl.

8. A compound as claimed in claim 7 wherein the group of formula (II) is an alkylamino or cycloalkylamino group preferably selected from optionally substituted methylamino, ethylamino, propylamino, isopropylamino, butylamino, cyclobutylamino, pentylamino, cyclopentylamino, hexylamino, 5 cyclohexylamino, heptylamino, cycloheptylamino, octylamino and cyclooctylamino.

9. A compound as claimed in any one of claims 1 to 8 wherein  $R^1$  is substituted with one or more of  $OR^{24}$ , halogen,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl,  $C_{1-6}$  haloalkyl,  $C_{1-6}$  alkylaryl,  $C_{1-6}$  alkylheterocyclyl,  $(CH_2)_nOR^{24}$ , 10  $(CH_2)_nNR^{24}_2$ ,  $SR^{24}$ ,  $NO_2$ ,  $CN$ ,  $NR^{24}_2$ ,  $CO_2R^{24}$ ,  $NR^{24}C(O)R^{24}$ ,  $NR^{24}S(O)_2R^{24}$ ,  $COR^{24}$ ,  $CONR^{24}_2$ ,  $S(O)_2R^{24}$ ,  $S(O)R^{24}$  or  $SO_2NR^{24}_2$ ; wherein  $R^{24}$  is hydrogen,  $C_{1-4}$  alkyl or  $C_{6-12}$  aryl preferably phenyl, or  $C_{5-12}$  heterocyclyl preferably pyridine, and  $n$  is 1, 2, 3, 4, 5 or 6;

15

wherein two  $R^{24}$  in  $NR^{24}_2$  may optionally form a partially saturated, unsaturated or fully saturated three to seven membered ring containing one to three heteroatoms, said ring is preferably independently substituted with one or more of halogen,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{1-12}$  haloalkyl,  $C_{3-12}$  carbocyclyl,  $C_{3-12}$  heterocyclyl,  $OR^{25}$ ,  $SR^{25}$ ,  $NO_2$ ,  $CN$ ,  $NR^{25}_2$ ,  $NR^{25}COR^{25}$ , 20  $NR^{25}CONR^{25}_2$ ,  $NR^{25}CO_2R^{25}$ ,  $CO_2R^{25}$ ,  $COR^{25}$ ,  $CONR^{25}_2$ ,  $S(O)_2R^{25}$ ,  $SONR^{25}_2$ ,  $S(O)R^{25}$ ,  $SO_2NR^{25}_2$ , or  $NR^{25}S(O)_2R^{25}$ ; and each saturated carbon in the optional ring is further optionally and independently substituted by  $=O$ ,  $=S$ ,  $NNR^{26}_2$ ,  $=N-OR^{26}$ ,  $=NNR^{26}COR^{26}$ ,  $=NNR^{26}CO_2R^{26}$ ,  $=NNSO_2R^{26}$ , or  $=NR^{26}$ ; 25 and each substitutable nitrogen atom is optionally substituted by  $R^{27}$ ,  $COR^{27}$ ,  $SO_2R^{27}$  or  $CO_2R^{27}$ ;

wherein  $R^{25}$  is hydrogen,  $C_{1-12}$  alkyl,  $C_{6-12}$  carbocyclyl or  $C_{5-12}$  heterocyclyl, optionally substituted by one or more of  $C_{1-6}$  alkyl, halogen,  $C_{1-6}$  haloalkyl,



OR<sup>28</sup>, SR<sup>28</sup>, NO<sub>2</sub>, CN, NR<sup>28</sup>R<sup>28</sup>, NR<sup>28</sup>COR<sup>28</sup>, NR<sup>28</sup>CONR<sup>28</sup>R<sup>28</sup>, NR<sup>28</sup>COR<sup>28</sup>,  
 NR<sup>28</sup>CO<sub>2</sub>R<sup>28</sup>, CO<sub>2</sub>R<sup>28</sup>, COR<sup>28</sup>, CONR<sup>28</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>28</sup>, SONR<sup>28</sup><sub>2</sub>, S(O)R<sup>28</sup>,  
 SO<sub>2</sub>NR<sup>28</sup>R<sup>28</sup>, NR<sup>28</sup>S(O)<sub>2</sub>R<sup>28</sup>, wherein the C<sub>1-12</sub> alkyl group optionally  
 incorporates one or two insertions selected from the group consisting of -O-, -  
 5 N(R<sup>28</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>28</sup> may be the same or different and  
 is as defined below;

wherein R<sup>26</sup> is hydrogen, C<sub>1-12</sub> alkyl, C<sub>6-12</sub> carbocyclyl or C<sub>5-12</sub> heterocyclyl,  
 optionally substituted by one or more of C<sub>1-6</sub> alkyl, halogen, C<sub>1-6</sub> haloalkyl,  
 10 OR<sup>28</sup>, SR<sup>28</sup>, NO<sub>2</sub>, CN, NR<sup>28</sup>R<sup>28</sup>, NR<sup>28</sup>COR<sup>28</sup>, NR<sup>28</sup>CONR<sup>28</sup>R<sup>28</sup>, NR<sup>28</sup>COR<sup>28</sup>,  
 NR<sup>28</sup>CO<sub>2</sub>R<sup>28</sup>, CO<sub>2</sub>R<sup>28</sup>, COR<sup>28</sup>, CONR<sup>28</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>28</sup>, S(O)R<sup>28</sup>, SO<sub>2</sub>NR<sup>28</sup>R<sup>28</sup>,  
 NR<sup>28</sup>S(O)<sub>2</sub>R<sup>28</sup>, wherein the C<sub>1-12</sub> alkyl group optionally incorporates one or two  
 insertions selected from the group consisting of -O-, -N(R<sup>28</sup>)-, -S(O)- and -  
 S(O<sub>2</sub>)-, wherein each R<sup>28</sup> may be the same or different and is as defined below;

15

wherein R<sup>27</sup> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl or C<sub>6-12</sub> aryl;

wherein R<sup>28</sup> is hydrogen, C<sub>1-6</sub> alkyl, or C<sub>1-6</sub> haloalkyl.

20 10. A compound as claimed in any one of claims 1 to 9 wherein R<sup>4</sup> is  
 selected from phenyl, cyclohexyl, acridine, benzimidazole, benzofuran,  
 benzothiophene, benzoxazole, benzothiazole, indole, isoindole, indolizine,  
 indazole, isoindole, isoquinoline, morpholine, naphthalene, phenazine,  
 phenothiazine, phenoxazine, piperazine, piperidine, pyridazine, pyridine,  
 25 pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, quinoline, quinolizine, tetrazine,  
 thiomorpholine, thianaphthalene, thiopyran, triazine, trithiane, furan, imidazole,  
 isoxazole, isothiazole, oxazole, oxadiazole, oxathiazole, pyrazole, pyrrole,  
 tetrazole, thiophene, thiadiazole, thiatriazole, thiazole or triazole, wherein each  
 substitutable carbon or hetero-atom in R<sup>4</sup> is optionally and independently



substituted by one or more of  $C_{1-6}$  alkyl,  $OR^{20}$ ,  $SR^{20}$ ,  $NO_2$ ,  $CN$ ,  $NR^{20}_2$ ,  $NR^{20}COR^{20}$ ,  $NR^{20}CONR^{20}_2$ ,  $NR^{20}COR^{20}$ ,  $NHCO_2R^{20}$ ,  $CO_2R^{20}$ ,  $COR^{20}$ ,  $CONR^{20}_2$ ,  $S(O)_2R^{20}$ ,  $SONR^{20}_2$ ,  $S(O)R^{20}$ ,  $SO_2NR^{20}_2$ , or  $NR^{20}S(O)_2R^{20}$ ;

5 wherein  $R^{20}$  is hydrogen,  $C_{1-6}$  alkyl, or  $C_{1-6}$  haloalkyl.

11. A compound as claimed in any one of claims 1 to 10 wherein  $R^4$  is a six-membered carbocyclyl or heterocyclyl group optionally substituted with one or more of  $OR^{29}$ ,  $NR^{29}_2$ ,  $SR^{29}$ ,  $(CH_2)_nOR^{29}$ ,  $(CH_2)_nNR^{29}_2$ , halogen,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl, haloalkyl,  $NO_2$ ,  $CN$ ,  $NR^{29}C(O)R^{29}$ ,  $NR^{29}S(O)_2R^{29}$ ,  $CO_2R^{29}$ ,  $COR^{29}$ ,  $CONR^{29}_2$ ,  $S(O)_2R^{29}$ ,  $S(O)R^{29}$  or  $SO_2NR^{29}_2$ ;

wherein  $R^{29}$  is hydrogen,  $C_{1-4}$  alkyl,  $C_{5-12}$  heterocyclyl or  $C_{6-12}$  aryl preferably phenyl, and  $n$  is 1, 2, 3, 4, 5 or 6;

15 wherein two  $R^{29}$  in  $NR^{29}_2$  may optionally form a partially saturated, unsaturated or fully saturated five to seven membered ring containing one to three heteroatoms, optionally and independently substituted with one or more of halogen,  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{1-12}$  haloalkyl,  $C_{6-12}$  carbocyclyl,  $C_{5-12}$  heterocyclyl,  $OR^{30}$ ,  $SR^{30}$ ,  $NO_2$ ,  $CN$ ,  $NR^{30}_2$ ,  $NR^{30}COR^{30}$ ,  $NR^{30}CONR^{30}_2$ ,  $NR^{30}COR^{30}$ ,  $NR^{30}CO_2R^{30}$ ,  $CO_2R^{30}$ ,  $COR^{30}$ ,  $CONR^{30}_2$ ,  $S(O)_2R^{30}$ ,  $SONR^{30}_2$ ,  $S(O)R^{30}$ ,  $SO_2NR^{30}_2$ , or  $NR^{30}S(O)_2R^{30}$ ; and each saturated carbon in the optional ring is further optionally and independently substituted by  $=O$ ,  $=S$ ,  $NNR^{31}_2$ ,  $=N-OR^{31}$ ,  $=NNR^{31}COR^{31}$ ,  $=NNR^{31}CO_2R^{31}$ ,  $=NNSO_2R^{31}$ , or  $=NR^{31}$ ; and each substitutable nitrogen atom is optionally substituted by  $R^{32}$ ,  $COR^{32}$ ,  $SO_2R^{32}$  or  $CO_2R^{32}$ ;

25

wherein  $R^{30}$  is hydrogen,  $C_{1-12}$  alkyl,  $C_{6-12}$  carbocyclyl or  $C_{5-12}$  heterocyclyl, optionally substituted by one or more of  $C_{1-6}$  alkyl, halogen,  $C_{1-6}$  haloalkyl,  $OR^{33}$ ,  $SR^{33}$ ,  $NO_2$ ,  $CN$ ,  $NR^{33}R^{33}$ ,  $NR^{33}COR^{33}$ ,  $NR^{33}CONR^{33}R^{33}$ ,  $NR^{33}COR^{33}$ ,  $NR^{33}CO_2R^{33}$ ,  $CO_2R^{33}$ ,  $COR^{33}$ ,  $CONR^{33}_2$ ,  $S(O)_2R^{33}$ ,  $SONR^{33}_2$ ,  $S(O)R^{33}$ ,

$\text{SO}_2\text{NR}^{33}\text{R}^{33}$ ,  $\text{NR}^{33}\text{S}(\text{O})_2\text{R}^{33}$ , wherein the  $\text{C}_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-\text{O}-$ ,  $-\text{N}(\text{R}^{33})-$ ,  $-\text{S}(\text{O})-$  and  $-\text{S}(\text{O}_2)-$ , wherein each  $\text{R}^{33}$  may be the same or different and is as defined below;

5


wherein  $\text{R}^{31}$  is hydrogen,  $\text{C}_{1-12}$  alkyl,  $\text{C}_{6-12}$  carbocyclyl or  $\text{C}_{5-12}$  heterocyclyl, optionally substituted by one or more of  $\text{C}_{1-6}$  alkyl, halogen,  $\text{C}_{1-6}$  haloalkyl,  $\text{OR}^{33}$ ,  $\text{SR}^{33}$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{NR}^{33}\text{R}^{33}$ ,  $\text{NR}^{33}\text{COR}^{33}$ ,  $\text{NR}^{33}\text{CONR}^{33}\text{R}^{33}$ ,  $\text{NR}^{33}\text{COR}^{33}$ ,  $\text{NR}^{33}\text{CO}_2\text{R}^{33}$ ,  $\text{CO}_2\text{R}^{33}$ ,  $\text{COR}^{33}$ ,  $\text{CONR}^{33}_2$ ,  $\text{S}(\text{O})_2\text{R}^{33}$ ,  $\text{S}(\text{O})\text{R}^{33}$ ,  $\text{SO}_2\text{NR}^{33}\text{R}^{33}$ ,  
 10  $\text{NR}^{33}\text{S}(\text{O})_2\text{R}^{33}$ , wherein the  $\text{C}_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-\text{O}-$ ,  $-\text{N}(\text{R}^{33})-$ ,  $-\text{S}(\text{O})-$  and  $-\text{S}(\text{O}_2)-$ , wherein each  $\text{R}^{21}$  may be the same or different and is as defined below;

wherein  $\text{R}^{32}$  is hydrogen,  $\text{C}_{6-12}$  aryl,  $\text{C}_{1-6}$  alkyl or  $\text{C}_{1-6}$  haloalkyl;

15

wherein  $\text{R}^{33}$  is hydrogen,  $\text{C}_{1-6}$  alkyl, or  $\text{C}_{1-6}$  haloalkyl.

12. A compound as claimed in any one of claims 1 to 11 wherein  $\text{R}^4$  is a five-membered heterocyclyl,



20

wherein A,  $\text{X}^2$ ,  $\text{Y}^2$  or Z are independently selected from N, O, C, S and M is C or N, wherein one, two, three or four of A,  $\text{X}^2$ ,  $\text{Y}^2$ , Z and M is other than C;

$\text{R}^{34}$ ,  $\text{R}^{35}$ ,  $\text{R}^{36}$  or  $\text{R}^{37}$  are independently selected from a lone electron pair,

25 hydrogen, halogen,  $\text{C}_{1-12}$  alkyl,  $\text{C}_{1-12}$  haloalkyl,  $\text{OR}^{38}$ ,  $\text{SR}^{38}$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{NR}^{38}_2$ ,  $\text{NR}^{38}\text{COR}^{38}$ ,  $\text{NR}^{38}\text{CONR}^{38}_2$ ,  $\text{NR}^{38}\text{COR}^{38}$ ,  $\text{NR}^{38}\text{CO}_2\text{R}^{38}$ ,  $(\text{CH}_2)_n\text{OR}^{38}$ ,

$(\text{CH}_2)_n\text{NR}^{38}_2$ ,  $\text{CO}_2\text{R}^{38}$ ,  $\text{COR}^{38}$ ,  $\text{CONR}^{38}_2$ ,  $\text{S}(\text{O})_2\text{R}^{38}$ ,  $\text{SONR}^{38}_2$ ,  $\text{S}(\text{O})\text{R}^{38}$ ,  $\text{SO}_2\text{NR}^{38}_2$ , or  $\text{NHS}(\text{O})_2\text{R}^{38}$ ;

wherein  $n$  is 1 to 6, preferably  $n$  is 1, 2 or 3;

- 5 or wherein any two of  $\text{R}^{34}$ ,  $\text{R}^{35}$ ,  $\text{R}^{36}$  or  $\text{R}^{37}$  may optionally form a partially saturated, unsaturated or fully saturated five to seven membered ring containing zero to three heteroatoms, each saturated carbon in the optional fused ring is further optionally and independently substituted with one or more of halogen,  $\text{C}_{1-12}$  alkyl,  $\text{C}_{2-12}$  alkenyl,  $\text{C}_{2-12}$  alkynyl,  $\text{C}_{1-12}$  haloalkyl,  $\text{C}_{6-12}$  carbocyclyl,  $\text{C}_{5-12}$  heterocyclyl,  $\text{OR}^{38}$ ,  $\text{SR}^{38}$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{NR}^{38}_2$ ,  $\text{NR}^{38}\text{CONR}^{38}_2$ ,  $\text{NR}^{38}\text{COR}^{38}$ ,  
 10  $\text{NR}^{38}\text{CO}_2\text{R}^{38}$ ,  $(\text{CH}_2)_n\text{OR}^{38}$ ,  $(\text{CH}_2)_n\text{NR}^{38}_2$ ,  $\text{CO}_2\text{R}^{38}$ ,  $\text{COR}^{38}$ ,  $\text{CONR}^{38}_2$ ,  $\text{S}(\text{O})_2\text{R}^{38}$ ,  $\text{SONR}^{38}_2$ ,  $\text{S}(\text{O})\text{R}^{38}$ ,  $\text{SO}_2\text{NR}^{38}_2$ , or  $\text{NR}^{38}\text{S}(\text{O})_2\text{R}^{38}$ ; and each saturated carbon in the optional fused ring is further optionally and independently substituted by  $=\text{O}$ ,  $=\text{S}$ ,  $\text{NNR}^{39}_2$ ,  $=\text{N}-\text{OR}^{39}$ ,  $=\text{NNR}^{39}\text{COR}^{39}$ ,  $=\text{NNR}^{39}\text{CO}_2\text{R}^{39}$ ,  $=\text{NNSO}_2\text{R}^{39}$ , or  
 15  $=\text{NR}^{39}$ ; and each substitutable nitrogen atom in  $\text{R}^4$  is optionally substituted by  $\text{R}^{40}$ ,  $\text{COR}^{40}$ ,  $\text{SO}_2\text{R}^{40}$  or  $\text{CO}_2\text{R}^{40}$ ;

wherein  $n$  is 1 to 6, preferably  $n$  is 1, 2 or 3;

- wherein  $\text{R}^{38}$  is hydrogen,  $\text{C}_{1-12}$  alkyl,  $\text{C}_{6-12}$  carbocyclyl or  $\text{C}_{5-12}$  heterocyclyl,  
 20 optionally substituted by one or more of  $\text{C}_{1-6}$  alkyl, halogen,  $\text{C}_{1-6}$  haloalkyl,  $\text{OR}^{41}$ ,  $\text{SR}^{41}$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{NR}^{41}\text{R}^{41}$ ,  $\text{NR}^{41}\text{CONR}^{41}\text{R}^{41}$ ,  $\text{NR}^{41}\text{COR}^{41}$ ,  $\text{NR}^{41}\text{CO}_2\text{R}^{41}$ ,  $\text{CO}_2\text{R}^{41}$ ,  $\text{COR}^{41}$ ,  $\text{CONR}^{41}_2$ ,  $\text{S}(\text{O})_2\text{R}^{41}$ ,  $\text{SONR}^{41}_2$ ,  $\text{S}(\text{O})\text{R}^{41}$ ,  $\text{SO}_2\text{NR}^{41}\text{R}^{41}$ ,  $\text{NR}^{41}\text{S}(\text{O})_2\text{R}^{41}$ , wherein the  $\text{C}_{1-12}$  alkyl group optionally incorporates one or two insertions selected from the group consisting of  $-\text{O}-$ ,  $-\text{N}(\text{R}^{41})-$ ,  $-\text{S}(\text{O})-$  and  $-\text{S}(\text{O}_2)-$ , wherein each  $\text{R}^{41}$  may be the same or different and is as defined below;  
 25

wherein  $\text{R}^{39}$  is hydrogen,  $\text{C}_{1-12}$  alkyl, carbocyclyl or heterocyclyl, optionally substituted by one or more of  $\text{C}_{1-6}$  alkyl, halogen,  $\text{C}_{1-6}$  haloalkyl,  $\text{OR}^{41}$ ,  $\text{SR}^{41}$ ,  $\text{NO}_2$ ,  $\text{CN}$ ,  $\text{NR}^{41}\text{R}^{41}$ ,  $\text{NR}^{41}\text{COR}^{41}$ ,  $\text{NR}^{41}\text{CONR}^{41}\text{R}^{41}$ ,  $\text{NR}^{41}\text{COR}^{41}$ ,  $\text{NR}^{41}\text{CO}_2\text{R}^{41}$ ,

CO<sub>2</sub>R<sup>41</sup>, COR<sup>41</sup>, CONR<sup>41</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>41</sup>, S(O)R<sup>41</sup>, SO<sub>2</sub>NR<sup>41</sup>R<sup>41</sup>, NR<sup>41</sup>S(O)<sub>2</sub>R<sup>41</sup>, wherein the C<sub>1-12</sub> alkyl group optionally incorporates one or two insertions selected from the group consisting of -O-, -N(R<sup>41</sup>)-, -S(O)- and -S(O<sub>2</sub>)-, wherein each R<sup>41</sup> may be the same or different and is as defined below;

5


























wherein R<sup>40</sup> is hydrogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl or C<sub>6-12</sub> aryl.


























wherein R<sup>41</sup> is hydrogen, C<sub>1-6</sub> alkyl, or C<sub>1-6</sub> haloalkyl.

10


13. A compound as claimed in claim 12 wherein R<sup>4</sup> is furan, imidazole, isoxazole, isothiazole, oxazole, oxadiazole, oxatriazole, pyrazole, pyrrole, tetrazole, thiophene, thiadiazole, thiatriazole, thiazole or triazole; and R<sup>34</sup>, R<sup>35</sup>, R<sup>36</sup> or R<sup>37</sup> are independently selected from a lone electron pair, hydrogen, halogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> haloalkyl, OR<sup>42</sup>, SR<sup>42</sup>, CN, NR<sup>42</sup><sub>2</sub>, NR<sup>42</sup>COR<sup>42</sup>, CO<sub>2</sub>R<sup>42</sup>, COR<sup>42</sup>, CONR<sup>42</sup><sub>2</sub>, S(O)<sub>2</sub>R<sup>42</sup>, or S(O)R<sup>42</sup>; wherein R<sup>42</sup> is hydrogen, C<sub>1-4</sub> alkyl, preferably methyl or ethyl or carbocyclyl, preferably phenyl.

20 14 A compound as claimed in any one of claims 1 to 13 selected from











15. A process for the manufacture of a compound of formula (I) wherein  $R^1$  is a group of formula (II) as defined in the any one of claims 1 to 14 of the invention comprising the condensation of an intermediate (III) with an intermediate (IV).



- wherein  $R^2$  and  $R^4$  are as defined in any one of claims 1 to 14; and wherein each of  $L^1$  and  $L^2$  is independently a leaving group wherein  $L^1$  and  $L^2$  together form a condensation product.

16. A process as claimed in claim 15 wherein  $L^1$  is OH,  $OR^{50}$ , OM, Cl, Br or I wherein  $R^{50}$  is  $C_{1-6}$  alkyl, preferably methyl or ethyl and M is Na, Li, K, Ca, Mg or Ba, and  $L^2$  is hydrogen or M.

17. A compound of formula (III)



(III)


wherein  $R^4$  is as defined any one of claims 1 to 14

$L^1$  is OH,  $OR^{50}$ , OM, Cl, Br, or I

$R^{50}$  is  $C_{1-6}$  alkyl, and

5 M is Na, Li, K, Ca, Mg, or Ba.

18. A process for the manufacture of a compound of formula (V) comprising removal of group  $R^{51}$  from an intermediate (VI)



(VI)

(V)

10

wherein  $L^3$  is  $R^1$  or  $L^1$ ;

$R^1$  and  $R^4$  are as defined in any one of claims 1 to 14;

$L^1$  is as defined in claim 17;

and  $R^{51}$  is an amino protecting group selected from  $R^{52}SO_2$ ,  $R^{52}C(O)$ ,  $R^{52}_3Si$ ,

15  $R^{52}OCH_2$ ,  $(R^{52})_2NSO_2$ ,  $(R^{52})_2NC(O)$ ,  $R^{52}OC(O)$ ,  $R^{52}(R^{52}O)CH$ ,  $R^{52}CH_2CH_2$ ,

$R^{52}CH_2$ ,  $PhC(O)CH_2$ ,  $CH_2=CH$ ,  $ClCH_2CH_2$ ,  $Ph_3C$ ,  $Ph_2(4\text{-pyridyl})C$ ,  $Me_2N$ ,


$HO-CH_2$ ,  $R^{52}OCH_2$ ,  $(R^{52})_3SiOCH_2$ ,  $(R^{52}O)_2CH$ ,  $t\text{-BuOC}(O)CH_2$ ,  $Me_2NCH_2$ ,

and tetrahydropyranylamine,

wherein  $R^{52}$  is  $C_{1-6}$  alkyl or  $C_{6-12}$  aryl.

20

19. A compound of formula (VI)




(VI)

wherein  $R^4$  is as defined in any one of claims 1 to 14, and

wherein  $L^3$  and  $R^{51}$  are as defined in claim 18.

- 5 20. A process for the manufacture of a compound of formula (VI) comprising a a) reaction of a compound of formula (VII) with stannane  $R^4$ - $\text{Sn}(R^{53})_3$  in the presence of a palladium catalyst or b) reaction of a compound of formula (VII) with boronic acid or ester  $R^4$ - $\text{B}(\text{OR}^{54})_2$  in a presence of a suitable palladium catalyst or c) reaction of a compound of formula (VII) with silane
- 10  $R^4$ - $\text{Si}(R^{55})_3$  in the presence of a palladium catalyst;



wherein  $R^4$  is as defined in any one of claims 1 to 14,

$L^3$  is as defined in claim 18;

$R^{51}$  is an amino protecting group as defined in claim 18;

- 15  $X^3$  is F, Cl, Br I or  $\text{CF}_3\text{SO}_3$ ,

and  $R^{53}$  is independently  $\text{C}_{1-6}$  alkyl;


- $R^{54}$  is independently hydrogen or  $\text{C}_{1-6}$  alkyl or wherein two  $R^{54}$  groups together optionally form a five, six or seven membered ring with the boron and oxygen atoms, wherein the ring is optionally substituted with one or more  $\text{C}_{1-6}$  alkyl group.
- 20

and  $R^{55}$  is independently  $\text{C}_{1-6}$  alkyl, F, or OH.

21. A process as claimed in claim 20 wherein the catalyst is one or more selected from  $(PPh_3)_2PdCl_2$ ,  $(PPh_3)_4Pd$ ,  $Pd(OAc)_2$ ,  $[PdCl(\eta^3-C_3H_5)]_2$ ,  $Pd_2(dba)_3$ ,  $Pd(dba)_2$  (dba = dibenzylidenacetone), and  $Pd/P(t-Bu)_3$ .

5

22. A compound of formula (VII)




(VII)

wherein  $L^3$  is as defined in claim 18;

wherein  $R^{51}$  is an amino protecting group as defined in claim 18;

10 wherein  $X^3$  is as defined in claim 20.

23. A process for the manufacture of a compound of formula (VII) comprising protection of the pyrrole nitrogen with a group  $R^{51}$ ,



(VIII)


(VII)

15 wherein  $L^3$  is as defined in claim 18;

wherein  $R^{51}$  is an amino protecting group defined in claim 18;

wherein  $X^3$  is as defined in claim 20.

24. A compound of formula (VIII)




(VIII)

wherein  $L^3$  is as defined in claim 18;

and  $X^3$  is as defined in claim 20.

- 5 25. A process for the production of a compound of formula (VIII) by the introduction of an  $X^3$  group into a compound of formula (IX)




(IX)

(VIII)


wherein  $L^3$  is as defined in claim 18 and  $X^3$  is as defined in claim 20.

- 10 26. A compound of formula (IX)



(IX)


wherein  $L^3$  is a group  $L^1$  as defined in claim 17 or a group  $R^1$ , wherein  $R^1$  is a group of formula (II)



(II)

- 15 wherein  $X$  is  $NR^3$ ,  $O$ ,  $S$  or  $(CR^{22}R^{22})_n$ ,  $Y$  is absent or is  $NR^{23}$ ,  $O$ , or  $(CR^{23}R^{23})_n$ ,  $R^2$  is optionally substituted  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{3-12}$  carbocyclyl or  $C_{3-12}$  heterocyclyl as claimed in any one of claims 1 to 14.

27. A process for the production of a compound of formula (VII) by the introduction of a  $X^3$  group to a compound of formula (X)




5 wherein  $L^3$  and  $R^{51}$  are as defined in claim 18 and  $X^3$  is as defined in claim 20.

28. A compound of formula (X)




10 wherein  $L^3$  is a group  $L^1$  as defined in claim 17 or a group  $R^1$ , wherein  $R^1$  is a group of formula (II)



15 wherein X is  $NR^3$ , O, S or  $(CR^{22}R^{22})_n$ , Y is absent or is  $NR^{23}$ , O, or  $(CR^{23}R^{23})_n$ ,  $R^2$  is optionally substituted  $C_{1-12}$  alkyl,  $C_{2-12}$  alkenyl,  $C_{2-12}$  alkynyl,  $C_{3-12}$  carbocyclyl or  $C_{3-12}$  heterocyclyl as claimed in any one of claims 1 to 14; and  $R^{51}$  is an amino protecting group as defined in claim 18.

29. A process for the preparation of compound of formula (IX) by the acid-catalysed hydrolysis of nitrile (XI) in the presence of an alcohol,







wherein  $L^3$  is  $OR^{50}$ ;

and  $R^{50}$  is as defined in claim 16.


5 30. A compound of formula (XI)



31. A process for the production of 1H-Pyrrolo[2,3-b]pyridine-5-carbonitrile (XI) comprising reaction of 5-bromo-1H-pyrrolo[2,3-b]pyridine with  $Zn(CN)_2$  in the presence of a palladium catalyst.



32 A compound as claimed in any one of claims 17, 19, 22, 24, 26, 28 or 30 selected from one or more of:



33. A process for the production of a compound of formula (I) comprising converting a starting material into an intermediate compound of any one of claims 17, 19, 22, 24, 26, 28 or 30, using a process as set out in one or more of claims 15, 16, 18, 20, 21, 23, 25, 27, 29 or 31.

and optionally converting the intermediate compound so formed into another intermediate compound

10

and then converting the intermediate compound into a compound of formula (I) using a process as claimed in any one of claims 15, 16, 18, 20, 21, 23, 25, 27, 29 or 31.

15 34. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 14 in combination with a pharmaceutically acceptable carrier, diluent or excipient.

35. A composition as claimed in claim 34, additionally comprising one or more of an anti-inflammatory agent, an AMPA receptor antagonist, a chemotherapeutic agent and/or an antiproliferative agent.
- 5 36. A process for the manufacture of a composition according to claim 34 or claim 35, comprising combining a compound according to any one of claims 1 to 14 of the invention with a pharmaceutically acceptable carrier or diluent and optionally with any one or more of additional agents of claim 35.
- 10 37. A compound as claimed in any one of claims 1 to 14, or a composition as claimed in claim 34 or claim 35, for use in medicine.
38. A compound as defined in any of claims 1-14, or a composition as defined in any of claims 34 or 35, for inhibiting JNK.
- 15 39. A compound as defined in any of claims 1-14, or a composition as defined in any of claims 34 or 35, for selectively inhibiting JNK3.
40. A compound as defined in any of claims 1-14, or a composition as  
20 defined in any of claims 34 or 35, for use in the prevention or treatment of a JNK-mediated disorder.
41. A compound or a composition as claimed in claim 40, wherein the  
25 disorder is a neurodegenerative disorder (including dementia), inflammatory disease, a disorder linked to apoptosis, particularly neuronal apoptosis, autoimmune disease, destructive bone disorder, proliferative disorder, cancer, infectious disease, allergy, ischemia reperfusion injury, heart attack, angiogenic disorder, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin

induced platelet aggregation and/or any condition associated with prostaglandin endoperoxidase synthase-2.

42. A compound or composition as claimed in claim 41, wherein the  
5 neurodegenerative disorder results from apoptosis and/or inflammation.

43. A compound or composition as claimed in claim 41 or 42, wherein the  
neurodegenerative disorder is: dementia; Alzheimer's disease; Parkinson's  
disease; Amyotrophic Lateral Sclerosis; Huntington's disease; senile chorea;  
10 Sydenham's chorea; hypoglycemia; head and spinal cord trauma including  
traumatic head injury; acute and chronic pain; epilepsy and seizures;  
olivopontocerebellar dementia; neuronal cell death; hypoxia-related  
neurodegeneration; acute hypoxia; glutamate toxicity including glutamate  
neurotoxicity; cerebral ischemia; dementia linked to meningitis and/or neurosis;  
15 cerebrovascular dementia; or dementia in an HIV-infected patient.

44. A compound or composition as claimed in claim 41 or 42, wherein the  
neurodegenerative disorder is a peripheral neuropathy, including  
mononeuropathy, multiple mononeuropathy or polyneuropathy, such as may be  
20 found in diabetes mellitus, Lyme disease or uremia; peripheral neuropathy  
caused by a toxic agent; demyelinating disease such as acute or chronic  
inflammatory polyneuropathy, leukodystrophies or Guillain-Barré syndrome;  
multiple mononeuropathy secondary to a collagen vascular disorder; multiple  
mononeuropathy secondary to sarcoidosis; multiple mononeuropathy secondary  
25 to a metabolic disease; or multiple mononeuropathy secondary to an infectious  
disease.

45. A compound or composition as claimed in claim 41, wherein the  
disorder is inflammatory bowel disorder; bronchitis; asthma; acute pancreatitis;

chronic pancreatitis; allergies of various types; Alzheimer's disease; autoimmune disease such as rheumatoid arthritis, systemic lupus erythematosus, glomerulonephritis, scleroderma, chronic thyroiditis, Graves's disease, autoimmune gastritis, diabetes, autoimmune haemolytic anaemia, autoimmune  
5 neutropaenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, ulcerative colitis, Crohn's disease, psoriasis or graft vs host disease.

46. A method of treating or preventing a JNK-mediated disorder in an  
10 individual, which method comprises administering to said individual a compound as claimed in any of claims 1-14 or a composition as claimed in any of claims 34 or 35.

47. A method as claimed in claim 46, wherein the individual is in need of  
15 the treatment or prevention of the disorder.

48. A method as claimed in claim 46 or 47, wherein the disorder is a neurodegenerative disorder, inflammatory disease, a disorder linked to apoptosis, particularly neuronal apoptosis, autoimmune disease, destructive  
20 bone disorder, proliferative disorder, cancer, infectious disease, allergy, ischemia reperfusion injury, heart attack, angiogenic disorder, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin induced platelet aggregation and/or any condition associated with prostaglandin endoperoxidase synthase-2.

25

49. A method as claimed in claim 48, wherein the neurodegenerative disorder results from apoptosis and/or inflammation.

50. A method as claimed in claim 48 or 49, wherein the neurodegenerative disorder is: dementia; Alzheimer's disease; Parkinson's disease; Amyotrophic Lateral Sclerosis; Huntington's disease; senile chorea; Sydenham's chorea; hypoglycemia; head and spinal cord trauma including traumatic head injury;  
5 acute and chronic pain; epilepsy and seizures; olivopontocerebellar dementia; neuronal cell death; hypoxia-related neurodegeneration; acute hypoxia; glutamate toxicity including glutamate neurotoxicity; cerebral ischemia; dementia linked to meningitis and/or neurosis; cerebrovascular dementia; or dementia in an HIV-infected patient.

10

51. A method as claimed in claim 48 or 49, wherein the neurodegenerative disorder is a peripheral neuropathy, including mononeuropathy, multiple mononeuropathy or polyneuropathy, such as may be found in diabetes mellitus, Lyme disease or uremia; peripheral neuropathy caused by a toxic agent;  
15 demyelinating disease such as acute or chronic inflammatory polyneuropathy, leukodystrophies or Guillain-Barré syndrome; multiple mononeuropathy secondary to a collagen vascular disorder; multiple mononeuropathy secondary to sarcoidosis; multiple mononeuropathy secondary to a metabolic disease, or multiple mononeuropathy secondary to an infectious disease.

20

52. A method as claimed in claim 46, 47 or 48, wherein the disorder is inflammatory bowel disorder; bronchitis; asthma; acute pancreatitis; chronic pancreatitis; allergies of various types; Alzheimer's disease; autoimmune disease such as rheumatoid arthritis, systemic lupus erythematosus,  
25 glomerulonephritis, scleroderma, chronic thyroiditis, Graves's disease, autoimmune gastritis, diabetes, autoimmune haemolytic anaemia, autoimmune neutropaenia, thrombocytopenia, atopic dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis, ulcerative colitis, Crohn's disease, psoriasis or graft vs host disease.



53 A method as claimed in any of claims 46-52, wherein one or more other active agent is administered to the individual simultaneously, subsequently or sequentially to administering the compound.

5

54. A method as claimed in claim 53, wherein the other active agent is an anti-inflammatory agent.

55. Use of a compound as defined in claim 1-14 in the manufacture of a medicament for the prevention or treatment of a JNK-mediated disorder.

10

56. Use as claimed in claim 55, wherein the disorder is a neurodegenerative disorder, inflammatory disease, a disorder linked to apoptosis, particularly neuronal apoptosis, autoimmune disease, destructive bone disorder, proliferative disorder, cancer, infectious disease, allergy, ischemia reperfusion injury, heart attack, angiogenic disorder, organ hypoxia, vascular hyperplasia, cardiac hypertrophy, thrombin induced platelet aggregation and/or any condition associated with prostaglandin endoperoxidase synthase-2.

15

57. Use as claimed in claim 55, wherein the neurodegenerative disorder results from apoptosis and/or inflammation.

20

58. Use as claimed in claim 56 or 57, wherein the neurodegenerative disorder is: dementia; Alzheimer's disease; Parkinson's disease; Amyotrophic Lateral Sclerosis; Huntington's disease; senile chorea; Sydenham's chorea; hypoglycemia; head and spinal cord trauma including traumatic head injury; acute and chronic pain; epilepsy and seizures; olivopontocerebellar dementia; neuronal cell death; hypoxia-related neurodegeneration; acute hypoxia; glutamate toxicity including glutamate neurotoxicity; cerebral ischemia;

25

dementia linked to meningitis and/or neurosis; cerebrovascular dementia; or dementia in an HIV-infected patient.

59. Use as claimed in claim 56 or 57, wherein the neurodegenerative  
5 disorder is a peripheral neuropathy, including mononeuropathy, multiple  
mononeuropathy or polyneuropathy, such as may be found in diabetes mellitus,  
Lyme disease or uremia; peripheral neuropathy caused by a toxic agent;  
demyelinating disease such as acute or chronic inflammatory polyneuropathy,  
leukodystrophies or Guillain-Barré syndrome; multiple mononeuropathy  
10 secondary to a collagen vascular disorder; multiple mononeuropathy secondary  
to sarcoidosis; multiple mononeuropathy secondary to a metabolic disease; or  
multiple mononeuropathy secondary to an infectious disease.

60. Use as claimed in claim 56, wherein the disorder is inflammatory bowel  
15 disorder; bronchitis; asthma; acute pancreatitis; chronic pancreatitis; allergies of  
various types; Alzheimer's disease; autoimmune disease such as rheumatoid  
arthritis, systemic lupus erythematosus, glomerulonephritis, scleroderma, chronic  
thyroiditis, Graves's disease, autoimmune gastritis, diabetes, autoimmune  
haemolytic anaemia, autoimmune neutropaenia, thrombocytopenia, atopic  
20 dermatitis, chronic active hepatitis, myasthenia gravis, multiple sclerosis,  
ulcerative colitis, Crohn's disease, psoriasis or graft vs host disease.

61. Use as claimed in any of claims 55-60, wherein the compound is  
administered simultaneously, subsequently or sequentially with one or more  
25 other active agent.

62. Use as claimed in claim 61, wherein the other active agent is an anti-  
inflammatory agent such as a p38 inhibitor.

63. An assay for determining the activity of the compounds as defined in any of claims 1-14, comprising providing a system for assaying the activity and assaying the activity of a compound as defined in any of claims 1-14.
- 5 64. An assay as claimed in claim 63, wherein the assay is for the JNK inhibiting activity of the compound, preferably for the JNK3-specific inhibiting activity of the compound.
65. An assay as claimed in claim 63 or 64, wherein the assay is a  
10 Scintillation Proximity Assay (SPA) using radiolabelled ATP, or is an ELISA.
66. A method of inhibiting the activity or function of a JNK, particularly JNK3, which method comprises exposing a JNK to a compound as defined in any of claims 1-14 or a composition as defined in any of claims 34-45.  
15
67. A method as claimed in claim 66, which is performed in a research model.
68. A method as claimed in claim 67, wherein the research model is an  
20 animal model.



